



Debunking the Myth of Exercise-Induced Immune Suppression: Redefining the Impact of Exercise on Immunological Health Across the Lifespan

John P. Campbell* and James E. Turner*

Department for Health, University of Bath, Bath, United Kingdom

OPEN ACCESS

Edited by:

Annemieke Boots,
University Medical Center
Groningen, Netherlands

Reviewed by:

Ana Maria Teixeira,
University of Coimbra, Portugal

Emily C. LaVoy,
University of Houston,
United States

*Correspondence:

John P. Campbell
j.campbell@bath.ac.uk;
James E. Turner
j.e.turner@bath.ac.uk

Specialty section:

This article was submitted
to Inflammation,
a section of the journal
Frontiers in Immunology

Received: 19 November 2017

Accepted: 15 March 2018

Published: 16 April 2018

Citation:

Campbell JP and Turner JE (2018)
Debunking the Myth of Exercise-
Induced Immune Suppression:
Redefining the Impact of Exercise
on Immunological Health
Across the Lifespan.
Front. Immunol. 9:648.
doi: 10.3389/fimmu.2018.00648

Epidemiological evidence indicates that regular physical activity and/or frequent structured exercise reduces the incidence of many chronic diseases in older age, including communicable diseases such as viral and bacterial infections, as well as non-communicable diseases such as cancer and chronic inflammatory disorders. Despite the apparent health benefits achieved by leading an active lifestyle, which imply that regular physical activity and frequent exercise enhance immune competency and regulation, the effect of a single bout of exercise on immune function remains a controversial topic. Indeed, to this day, it is perceived by many that a vigorous bout of exercise can temporarily suppress immune function. In the first part of this review, we deconstruct the key pillars which lay the foundation to this theory—referred to as the “open window” hypothesis—and highlight that: (i) limited reliable evidence exists to support the claim that vigorous exercise heightens risk of opportunistic infections; (ii) purported changes to mucosal immunity, namely salivary IgA levels, after exercise do not signpost a period of immune suppression; and (iii) the dramatic reductions to lymphocyte numbers and function 1–2 h after exercise reflects a transient and time-dependent redistribution of immune cells to peripheral tissues, resulting in a heightened state of immune surveillance and immune regulation, as opposed to immune suppression. In the second part of this review, we provide evidence that frequent exercise enhances—rather than suppresses—immune competency, and highlight key findings from human vaccination studies which show heightened responses to bacterial and viral antigens following bouts of exercise. Finally, in the third part of this review, we highlight that regular physical activity and frequent exercise might limit or delay aging of the immune system, providing further evidence that exercise is beneficial for immunological health. In summary, the over-arching aim of this review is to rebalance opinion over the perceived relationships between exercise and immune function. We emphasize that it is a misconception to label any form of acute exercise as immunosuppressive, and, instead, exercise most likely improves immune competency across the lifespan.

Keywords: exercise, physical activity, upper respiratory tract infections, open window hypothesis, infection susceptibility, ageing, immunosenescence, immune competency

INTRODUCTION

Lifelong physical activity¹ is a potent means of reducing the risk of non-communicable diseases including cancer, cardiovascular disease, and other chronic inflammatory disorders (1). Evidence also shows that a physically active lifestyle diminishes the risk of contracting a range of communicable diseases including viral and bacterial infections (2–6). In contrast to the widely accepted long-term health benefits that are achieved by regular physical activity, which imply that immune competency and regulation are improved by frequent exercise bouts, the effect of a single bout of exercise on immune function remains hotly disputed. Undeniably, acute vigorous exercise has a profound effect on the phenotypic makeup and functional capacity of the immune system. Indeed, the behavior of almost all immune cell populations in the bloodstream is altered in some way during and after exercise (7, 8). However, for decades, it has been widely accepted that these changes result in a temporary decline in immune competency in the hours following exercise. In the first part of this review, we re-interpret these data and strive to dispel the misconception that an acute bout of exercise is detrimental to immunological health. In the second part of this article, we demonstrate that rather than suppressing immunity, contemporary evidence shows that an acute bout of exercise improves immune surveillance, for example leading to enhanced antibacterial and antiviral immunity. Finally, in the third part of this article, we summarize recent data suggesting that regular physical activity and frequent exercise, which reduces systemic inflammatory activity and improves aspects of immune function, also leads to alterations in classical biomarkers of an aging immune system. These changes could be interpreted as limiting or delaying immunological aging (9–13).

PART A: IS IT TIME TO CLOSE THE SHUTTERS ON THE “OPEN-WINDOW” HYPOTHESIS? A BOUT OF EXERCISE DOES NOT SUPPRESS IMMUNE COMPETENCY

A prevailing myth has formed in the literature that participating in an acute bout of aerobic exercise, particularly if it is vigorous and prolonged, can be detrimental to immune competency. The foundations of this belief lay in research publications emanating from

the 1980s and 1990s, reviewed extensively elsewhere (7, 8, 14). Findings from these early studies led to three principles of exercise immunology being formed, which have, until now, generally been unchallenged in the literature: (i) infection risk is increased after an acute bout of prolonged and vigorous aerobic exercise; (ii) acute bouts of vigorous exercise can lead to a temporary reduction to salivary IgA levels culminating in a higher risk of opportunistic infections; and (iii) transient decreases in the number of peripheral blood immune cells, which occurs in the hours following vigorous exercise, represents a period of immune suppression. Over the years, these collective observations have coalesced and led to the creation of the so-called “open-window” hypothesis, which purports that the immune system is compromised in the hours after vigorous exercise, leading to an increased risk of opportunistic infections in the days thereafter. To this day, the “open window” hypothesis continues to be discussed (15), despite the existence of contradictory evidence. Here, in the first part of this review, we aim to dispel the “open-window” hypothesis by revisiting key research studies and highlighting that limited evidence exists to support each of the three pillars that lay its foundation.

Exercise and Opportunistic Infections

It has been speculated for over a century that participation in physical activity heightens the risk of opportunistic infections (16). However, deeper investigation into the relationship between exercise and infection susceptibility did not take place until the end of the twentieth century. At this juncture, the principle focus of many of these studies in the late 1980s and early 1990s was to determine whether infection incidence was increased in elite and recreational athletes in the weeks following mass participation distance running events. One of the first studies from this era found that one third of 150 runners participating in the 1982 Two Oceans 56 km ultramarathon in Cape Town South Africa self-reported symptoms of upper respiratory tract infections (URTIs; symptoms = runny nose, sore throat, sneezing) within 2 weeks of the race (17). The control group, who were age matched and shared a home with another of the race competitors, reported only half the amount of URTIs in the same period (17). Similar observations were made in an often-cited larger study of the 1987 Los Angeles Marathon (18). Of 2,311 respondents who had completed the marathon and whom did not report an infection in the week prior to the race, 12.9% reported an infection in the week after the race compared to only 2.2% of individuals who withdrew from the race for reasons other than illness (odds ratio of infection in runners versus non-runners = 5.9). A separate study conducted around the same time found that shorter duration running events, such as 5, 10 km, and half-marathons (21 km) did not appear to elicit an increased incidence of self-reported URTIs (19), thereby suggesting that URTI symptoms are increased only when the duration of the exercise is long.

A fundamental limitation of each of the aforementioned studies is that none of the self-reported infections were clinically confirmed by laboratory analyses (e.g., molecular or microbiological techniques, such as polymerase chain reaction or bacterial cultures). As a consequence, it was questioned whether the

¹“Physical activity” refers to activities undertaken during leisure time, at home, as part of employment, or for transport purposes. “Exercise” is a component of physical activity within the leisure time domain and refers to physical activities that are planned, structured, repetitive, and undertaken for the purpose of improving or maintaining components of physical fitness and/or sporting performance. When individuals are referred to as being “active” or “inactive,” the description infers that these people undertake (or fail to undertake) a defined level of exercise or physical activity (e.g., age-specific physical activity recommendations, such as those published by the World Health Organisation). In this review, the term “exercise” will generally be used to describe the effects that active behaviours have on immune competency. Individuals described as being “sedentary” accumulate prolonged periods of behaviour eliciting low energy expenditure (e.g. sitting and lying).

self-reported URTIs in these studies represented genuine infections. Clarifying this issue, a study employing nasopharyngeal and throat swabs in athletes who reported URTI symptoms over a 5-month period—including periods of competition—found that few of the self-reported infections were of bacterial, viral, chlamydial, or mycoplasmal nature (20). Indeed, of 37 episodes of URTI reported by athletes in this study, only 11 of these (30%) had a positive laboratory diagnosis. These findings place the previously discussed marathon studies in a different light (17, 18) and open the possibility that many of the URTIs reported were a symptom of other non-infectious causes. Indeed, of the of the non-infectious “URTIs” reported by Spence et al. (20), and likely captured elsewhere (17, 18), it is proposed that these symptoms are a result of other causes, including allergy and asthma, non-specific mucosal inflammation, or airway epithelial cell trauma due to increased ventilation or exposure to cold air (21). In the few cases of clinically confirmed URTIs, these appear to be from viruses—in particular rhinoviruses (i.e., the “common cold”)—rather than bacterial infection (20, 22), which is in line with the typical incidence and etiology of infections at the population level (23).

In the previously discussed clinical laboratory study of infection incidence in athletes (20), it is notable that the proportion of athletes with a confirmed infection during the 5-month period was as follows: 2/20 controls (10%), 3/31 (10%) recreational athletes, and 6/32 (19%) elite athletes. Although this study was small, these data appear to align with observations from earlier self-report studies which found that the incidence of URTI symptoms was higher in those with the fastest race time and those who had completed a greater training volume pre-race (17, 18). These early observations contributed to the formulation of the “J-shaped curve” (24). This hypothesis infers that those who undertake an excessive volume of exercise, over a period of weeks and months, sometimes referred to as “over-training” or “intensified training” (25, 26), are at a greater risk of infections (24). In this scenario, other factors present *prior* to an acute bout of exercise, such as psychological stress and anxiety (27–29), or nutritional deficiencies (30) which are known to impact immune regulation, are likely to impact immune competency and contribute to the risk of genuine URTIs, rather than the acute and transient immune changes that arise *after* the acute bout of exercise itself; these acute immunological changes arising after acute exercise are discussed later in this article (see Part A: “Is it Time to Close the Shutters on the “Open-Window” Hypothesis? A Bout of Exercise Does Not Suppress Immune Competency”; and see “Exercise and Salivary IgA and Changes to Lymphocyte Frequency and Functional Capacity in the Hours After Acute Exercise”).

Moreover, we contend that attendance at any mass participation event—whether it is a marathon or otherwise—is likely to increase the risk of acquiring novel infectious pathogens, which are in abundance due to the mass gathering of people. For example, it has been shown that around 40% of individuals attending the Hajj—a crowded religious event in Saudi Arabia—self-report an URTI (31). In this study, there was a greater risk of infection among those with the longest exposure to crowds (31). Thus, it is important to consider that other underlying factors, often not measured in the context of exercise and illness studies, likely

play a greater role in infection risk than exercise participation *per se*. For example, recently, it has been demonstrated that air travel is a significant predictor of illness symptoms in athletes (32). Infections linked to air travel are exacerbated by long-haul flights crossing multiple time zones, implicating many other factors known to influence immune function, including exposure to hypobaric hypoxia, radiation, temperature changes, sleep disruption, fatigue, altered or inadequate diet, dehydration, and psychological stress (32–34).

Contrary to the aforementioned reports that exercise heightens infection incidence, it is often overlooked that other studies indicate that exercise participation may in fact *reduce* the incidence of infections. For example, a recent prospective cohort study of 1,509 Swedish men and women aged 20–60 years found that higher physical activity levels were associated with a lower incidence of self-reported URTIs (35). A much smaller but very detailed analysis of illness records kept by 11 elite endurance athletes over a period of 3–16 years showed that the total number of training hours per year was inversely correlated with sickness days reported (36). Similarly, another study of swimmers monitored for 4 years found that national level athletes had higher incidence of infections than more elite international level athletes (37). Finally, studies of ultramarathon runners, who undertake the largest volume of exercise among athletes, have shown that these individuals report fewer days missed from school or work due to illness compared to the general population. For example, the mean number of sickness days reported over 12 months was 1.5 days in a study of 1,212 ultramarathon runners and 2.8 days in a study of 489 ultramarathon runners (38, 39). These studies compared their findings to data from the United States Department of Health and Human Services report in 2009, showing that the general population report on average 4.4 illness days each year. Thus, a number of studies challenge the “J-shaped curve,” indicating that athletes undertaking the largest training loads, become ill less frequently than athletes competing at, and training at, a lower level. These findings have previously been conceptualized by extending the “J-shaped curve” into an “S-shaped curve,” thereby suggesting that very elite athletes are better adapted to the demands of their training (40). Given the nature of their design, very few of these reports—akin to many of the aforementioned studies showing increased infection risk among athletes following mass participation endurance events—used appropriate laboratory diagnostics to confirm an infection. However, despite their limitations, it is important to highlight that there are as many epidemiological studies showing that regular exercise *reduces* infections as there are studies showing exercise *increases* infections, and that these studies are often overlooked in the exercise immunology literature.

It should also be considered whether the commonly reported “increased frequency” of illness symptoms among athletic populations or those taking part in sporting events is indeed more frequent than among the general population. For example, large studies have reported that approximately 7–10% of athletes competing in the Olympic Games report symptoms of illness during the competition weeks (41, 42). However, accumulating evidence suggests the incidence of infection among athletes is

not substantially different to other populations. For example, in a telephone survey of 2011 adults considered to represent the general population in the USA, 24% experienced a cold during a 4-week period, which is a similar timeframe to many international sporting competitions (43). In another telephone survey of 4,051 adults in the USA, 72% experienced at least 1 non-influenza related URTI over 12 months, and on average, experienced 2 infections annually (44). In a year-long Internet-based monitoring study of 627 individuals over 14 years of age in Germany, weekly acute respiratory illness rates were 2.7–8.2%, manifesting in 1.3–3.2 episodes annually (45). Thus, evidence suggests that the frequency of illness episodes in athletic communities is similar to the general population annually.

In the context of exercise participation in older age, it would appear to be counterintuitive that the incidence of URTI symptoms appears to be inversely correlated with age: it has been shown that URTI symptoms are more common in younger rather than older runners (18). While, once more, this aforementioned study did not confirm infections by laboratory analyses, it is notable that if acute exercise does suppress immune competency, it might be expected that older adults—whom typically have inferior immune function—would be at greatest risk of exercise-induced immune suppression. Rather than exercise *per se*, again, a more likely explanation for differences in illness symptoms between groups of athletes are other factors present before competing, such as fatigue, nutritional deficiency, psychological stress, or environmental exposures. On the other hand, proponents of the “open-window” hypothesis could portend that experienced athletes have higher tolerance of the symptoms associated with URTI, and/or have developed coping methods or strategies to reduce symptoms. Alternatively, experienced athletes may have evolved strategies to reduce infection risk by adopting good practice (e.g., sleep, diet, hygiene) before, during and following attendance at a mass participation event.

Separately, it has also been questioned whether illness symptoms—even if confirmed to be infectious—are a result of encountering a novel pathogen. For example, over a decade ago, some studies suggested that reactivation of latent viruses—such as *Epstein Barr Virus* which had most likely infected the host during childhood—was responsible for illness symptoms after exercise (46, 47). Although these studies suggest new pathogens were not to blame, it was interpreted that exercise-induced immune suppression had resulted in loss of viral control. However, herpes viruses can reactivate even with a fully functioning immune system, for example, in response to adrenergic activity, reactive oxygen species and inflammatory cytokines (48–50), all of which increase during exercise. Moreover, recent evidence appears to discount herpes virus involvement in causing symptoms of illness, by showing that individuals previously infected with *Cytomegalovirus*, or those infected with both *Epstein Barr Virus* and *Cytomegalovirus*, exhibited a lower incidence of illness symptoms than individuals not latently infected (51).

Taken together, evidence that participation in an acute bout of vigorous exercise leads to heightened infection incidence remains spurious. If symptoms of URTI are observed after a bout of vigorous exercise, the cause is unlikely to be infectious. However, if

infection or immune impairment is confirmed, their trigger is more likely to be the physical, nutritional, and psychological wellbeing of the individual prior to undertaking the single bout of acute vigorous exercise. In the context of mass participation sporting events, it is likely that increased exposure to pathogens, or the influence of environmental factors that can affect immune function (e.g., travel, sleep disruption) most likely explain genuine infections. Thus, we conclude that it is unlikely that vigorous and prolonged exercise heighten the risk of infections and should not be considered a deterrent to those seeking to become more physically active.

Exercise and Salivary IgA

A second mainstay of exercise immunology that has received considerable attention over the last three decades is the assessment of exercise-induced changes to mucosal immunity, principally *via* measurement of IgA levels in saliva (52). Given that IgA is the most abundant immunoglobulin in mucosal secretions and that its principle role is the inhibition of invading pathogens, isolated changes to salivary IgA following exercise has been considered of some importance in light of the purportedly higher risk of infections among athletes (7, 8).

One of the earliest and most cited papers in this research area found that IgA was reduced by 20% after 2–3 h of cross-country skiing (53). Another study found that this effect is transient, whereby salivary IgA concentrations decreased immediately after 2 h of intensive cycling exercise, and remained low in samples collected 1 h post-exercise, but returned to normal levels within 24 h (54). A criticism of these studies at that time was that the absolute IgA levels reported did not adequately control for the amount of saliva produced, and thus these results may misrepresent IgA secretion. Although some studies measuring IgA secretory rate (IgA protein concentration multiplied by saliva flow rate) support the early findings with IgA concentration, others have shown profoundly contradictory results. For example, in alignment with prior observations, it was found in trained runners that IgA secretion rate decreased by 25% from pre-marathon to 90 min post-marathon (55). Likewise, in a separate study, a 20% reduction in IgA secretion rate was observed in elite athletes after a 2-h rowing exercise session (56). Several other studies of similar design reported analogous findings (57–59); however, contradictory findings are also in abundance but are much less cited in the literature. Indeed, an elegant study exploring the effects of different exercise intensities, including moderate- and high-intensity exercise to exhaustion, found that although saliva flow rate decreased, IgA secretion rate actually increased in response to both of the exercise bouts. In the words of the authors, exercise to exhaustion has an “*effect on the quantity of saliva, but not the quality of saliva*” (60). Many other studies have also reported that exercise does not elicit a decrease in IgA secretion rates following exercise (61–66).

Any subtle isolated changes to IgA that occur after exercise appear to be clinically insignificant as it would appear that the increased incidence of URTI symptoms that has been purported following vigorous and prolonged exercise is unrelated to salivary immunoglobulin status. A longitudinal field study of participants in the Comrades Marathon (86.5 km ultramarathon) in South

Africa found that salivary IgA levels in the 4 weeks prior to the race, and 2 weeks following the race, were unrelated to the incidence of self-reported URTI (67). In the aforementioned study, it was found that symptoms of URTI were highest 4 weeks prior to the marathon, and URTI symptoms seemed to re-appear within many of the same athletes 1 or 2 weeks post-marathon. This study did not confirm the re-emergence of infections by laboratory testing, but if this was indeed demonstrated in future studies, it again shows that acute exercise participation *per se* does not heighten risk of opportunistic infections. In this case, an underlying infection, not resolved prior to exercise participation, or some other idiosyncrasy, is perhaps to blame. Such conclusions appear to be supported by evidence from athletes who report the most frequent illness symptoms. For example, these individuals exhibit mostly anti-inflammatory cytokine responses when whole blood, collected at rest, is cultured *ex vivo* with antigens from diphtheria, tetanus, acellular pertussis, poliomyelitis, and hemophilus influenza type b (68, 69). These findings suggest the immune system may be functionally altered by underlying illness, or is already different in these “illness-prone” athletes prior to infection, rather than exercise affecting immune function *per se*.

A flaw in studies investigating the link between mucosal immunity and purported exercise-induced infection risk is that oral health status is rarely adequately evaluated. Salivary IgA is heavily involved in host-bacterial ecology and mucosal homeostasis (70, 71). As optimal oral health is rare in adults—with nearly all exhibiting caries, gingivitis or periodontitis—profound between-person IgA variation has been reported, which is dependent on oral health status (71). Moreover, periodontal diseases are complex and multifactorial, and as a result, studies report large fluctuations in IgA levels relative to disease status between persons, probably due to the bespoke ecological makeup of different host mucosa (71). In addition, oral disease is a common problem in athlete populations (72), which is likely to be caused by high volume and frequent carbohydrate consumption, and in some, a neglect of oral hygiene, perhaps due to practical constraints. Thus, changes to oral inflammatory status has not been adequately considered, and emerging salivary biomarkers of oral inflammation, such as immunoglobulin-free light chains (73), may offer a means of controlling for this confounder. As highlighted elsewhere (70), salivary IgA is also highly vulnerable to short-term variation, in particular, due to circadian rhythms, typically peaking in the morning, and falling thereafter (74). As salivary IgA secretion is controlled by the parasympathetic and sympathetic nervous system, psychological stress also plays a powerful role in regulating IgA levels (75). Animal models suggest that salivary IgA levels could vary up to 27-fold within the same host over a short period of time (76). Salivary IgA levels are also affected by factors such as sex differences (77), diet, ethnicity, disease, medications, tobacco, and phase of the menstrual cycle, as reviewed elsewhere (71). To overcome some of these variations in salivary IgA, it is often the case that studies evaluate only secretory IgA (sIgA; i.e., IgA containing the secretory component) as this represents IgA produced by local mucosal plasma cells, and not IgA from the bloodstream transported *via* crevicular fluid. While this approach may reduce some confounding error, most IgA in saliva contains the secretory component and is, itself, subject to large variation;

extensively reviewed elsewhere (78). Given these many considerations, we propose that longitudinal measurement of salivary IgA, as an isolated measure of immune competency within a single host, and even more so between persons, depicts too confusing a picture, and it is ambitious to say that any subtle changes to salivary IgA following exercise reflects immune suppression and a heightened risk of opportunistic infections. Given the limitations of salivary IgA measurement, research is being undertaken to explore mucosal IgA in other biofluids, and a recent study has shown links between reduced tear IgA levels and infection incidence (79). Others have moved toward more comprehensive oral immunity panels (80), and such strategies could benefit further from an integrative approach that, in addition to immune parameters, incorporates full dental examination, oral inflammation biomarkers, and host mucosal ecology.

Changes to Lymphocyte Frequency and Functional Capacity in the Hours After Acute Exercise

One of the most reproduced findings in human exercise physiology is the profound and transient time-dependent change that arises to the phenotypic composition and functional capacity of lymphocytes in the peripheral bloodstream in response to a single bout of exercise (8). During vigorous aerobic exercise, it is commonly observed that peripheral blood lymphocyte frequency—and, concomitantly, the functional capacity of the lymphocyte pool—is dramatically increased, leading to the concept that, during exercise, exercise appears to “stimulate” the immune system. On the other hand, in the hours following exercise, it is typically observed that total peripheral blood lymphocyte frequency—and the functional capacity of the lymphocyte pool—is decreased below pre-exercise levels, leading some to propose that exercise induces a short-term window of immune suppression (termed the “open-window” hypothesis). The purpose of this part of our review is to outline that it is a misconception to state that the “reductions” to lymphocyte frequency and function, that arise in the hours following acute exercise, reflects immune suppression, and instead we emphasize that during this post-exercise period, the immune system is in a heightened state of immune surveillance and regulation.

Transient Changes to Blood Lymphocyte Frequency in the Hours Following Exercise

The classic biphasic response of lymphocytes to acute steady state vigorous exercise lasting for around at least 45–60 minutes, is first characterized by a dramatic lymphocytosis. This response is typified by a dramatic influx of natural killer cells, which rise by up to 10-fold, and CD8⁺ T cells which increase to a lesser—but still profound—extent by approximately 2.5-fold (81). This exercise intensity-dependent mobilization is driven in part by increased shear forces and blood pressure during exercise causing a non-specific flushing of the marginal pools (82) but, moreover, is principally governed by adrenergic stimulation of beta-2-adrenergic receptors on the surface of lymphocytes, arising from adrenaline released during exercise, causing endothelial detachment and subsequent recirculation

of lymphocytes into the bloodstream (83–85). Indeed, the lymphocyte mobilization response observed during exercise appears to broadly mirror the differential expression of beta-2 adrenergic receptors on lymphocytes: natural killer cells > CD8⁺ T cells > B cells > CD4⁺ T cells, including regulatory T cells (81, 86–88). Upon exercise cessation, the classic biphasic exercise response is next characterized by a dramatic decrease in the frequency of lymphocytes in the bloodstream. This nadir is typically observed approximately 1–2 h post-exercise when the lymphocyte numerical count is lower than pre-exercise levels; lymphocyte frequency normally returns to pre-exercise levels within 24 h (87, 89, 90). The lymphopenia that occurs 1–2 h later is exercise intensity dependent and the most profound reductions during this period are typically observed among natural killer cells and CD8⁺ T cells (90). Akin to purported reductions to salivary IgA, discussed earlier, it was perceived that the numerical decline of blood lymphocytes that arises during this time represented “double jeopardy” (89), temporarily exposing an individual to impaired immune competency and concomitantly providing an “open-window” for opportunistic infections (91, 92).

Rather than suppressing immune competency however, a more contemporary viewpoint is that this acute and transient lymphopenia 1–2 h after exercise is beneficial to immune surveillance and regulation. Indeed, in what appears to be a highly specialized and systematic response, it is widely proposed that exercise redeploys immune cells to peripheral tissues (e.g., mucosal surfaces) to conduct immune surveillance. Here, these immune cells are thought to identify and eradicate other cells infected with pathogens, or those that have become damaged or malignant, termed the acute stress/exercise immune-enhancement hypothesis (93). A seminal study by Kruger and colleagues, using fluorescent cell tracking in rodents, found that T cells are redeployed in large numbers to peripheral tissues including the gut and lungs, and to the bone marrow following exercise (84, 94). In line with Dhabhar’s theory, it is hypothesized that this redistribution reflects heightened immune surveillance in sites where pathogens are likely to be encountered during and after exercise (i.e., lungs, gut). This response has also been proposed to maintain immune homeostasis *via* augmented regulatory activities (12). In this context, evidence implies that bone marrow homing and subsequent apoptosis of senescent T cells stimulates the production or mobilization of new progenitor cells into the periphery (95), which has been hypothesized as an exercise-induced means of maintaining a younger immune system (12), discussed later in Part C “Does Exercise and Regular Physical Activity Influence Immunological Ageing.” Links between exercise-induced apoptosis and lymphopenia have in the past been interpreted as detrimental, with speculation that apoptosis could be responsible for the fall in lymphocyte number in the hours after exercise (96, 97). Other studies have reported increased lymphocyte apoptosis immediately after exercise (i.e., as a result of the large mobilization of cells) but not in the hours following exercise during lymphopenia (98–100). Although the magnitude of lymphocyte apoptosis reported in studies is dependent on the measurement technique, typically <10% of lymphocytes undergo post-exercise apoptosis (96, 97). Given the 30–60% decrease in lymphocyte

numbers post-exercise (89, 101, 102), apoptosis could be a small contributor to exercise-induced lymphopenia, but this process of cell death is likely to be beneficial given the stimulation of progenitor cells from the bone marrow (95).

While it has not been shown in humans that exercise—in line with rodent models—causes the redistribution of immune cells to peripheral tissues, further support for a coordinated, exercise-induced immune surveillance response elicited by lymphopenia, is revealed by studying the phenotypic characteristics of the cells that preferentially mobilize and subsequently extravasate out of the circulation after exercise. With regard to natural killer cells—the most exercise-responsive lymphocyte subset—CD56^{dim} cells are preferentially redeployed rather than their CD56^{bright} counterparts (81). CD56^{dim} cells are a mature subset of natural killer cells which have exclusive migratory potential for non-lymphoid tissue and potent effector capabilities, including the capacity to produce high amounts of perforin and granzyme, whereas CD56^{bright} cells are a more immature regulatory cell subset (103) and reside in secondary lymphoid organs, typified by their cell-surface expression of CD62L and CCR7 (104). CD56^{dim} natural killer cells can be further dissected, into cells with highly potent effector function based on loss of NKG2A and expression of killer immunoglobulin-like receptors and CD57 (105, 106). In a recent study, it was shown that these natural killer cells, which are capable of rapid effector functions, are preferentially redistributed after exercise (107, 108). Synergistically, T cells also appear to exert heterogeneous but highly coordinated responses to acute exercise. Indeed, it is consistently observed that discrete populations of CD8⁺ but not CD4⁺ T cell subsets are redeployed by exercise. For some time, there was confusion pertaining to the exact behavior of CD8⁺ T cells in response to exercise. Rather than losing or gaining markers of adhesion or activation—as evaluated elsewhere (8)—these changes represent a uniform redeployment of a preferentially mobilized group of memory cells. In a flurry of studies about a decade ago, it was shown that exercise selectively mobilizes memory CD8⁺ T cells with a phenotypic propensity for homing to peripheral tissues—typified for example by CD11, and not CCR7 or CD62L expression—and the distinctive capacity to mount rapid effector functions (81, 109–111). This response presumably facilitates the detection and elimination of neoplastic, stressed or infected cells in synergy with natural killer cells, as proposed elsewhere (112). Aligned with the immune surveillance theory of Burnet and Thomas, reviewed elsewhere (113), these results imply that sentinel cells of the immune system are redeployed by exercise-induced perturbations to stress hormones, to exert effector functions against neoplastic, stressed, or infected cells in the hours following exercise. This process, which occurs daily in a natural diurnal process (114), orchestrated subtly by stress hormones (115, 116), appears to be primed in response to exercise, leading to enhanced immune surveillance (117). Principles of the acute stress/exercise immune-enhancement hypothesis continue to be investigated. For example, it has recently been shown that acute exercise does not preferentially mobilize CD8⁺ T cells and natural killer cells with the capacity

for skin-homing (118). However, skin-homing in this context is a role that may be fulfilled by exercise-responsive mucosal-associated invariant T cells (119); but further work is needed in this area.

A key example illustrating how exercise-induced immune cell redistribution is beneficial to host health can be found in the rapidly emerging field of exercise oncology. Indeed, a recent seminal study demonstrated inhibition of tumor onset and disease progression across a range of tumor models in voluntarily active rodents (112). In this work, natural killer cell infiltration was significantly increased in tumors from active versus inactive rodents, leading to the conclusion that the presence of natural killer cells (but perhaps also T cells) in tumor sites, redeployed by adrenaline during exercise stress, “provides a spark” for tumor elimination, in what could be considered a form of “exercise immunotherapy” (112, 120). Importantly, administration of propranolol—a beta 2 adrenergic blocker—abolished the adrenaline-induced redistribution of immune cells, and nullified the anti-tumor effect of exercise on neoplastic growth (112). While these studies are limited to rodents, there is growing evidence that exercise may promote anti-cancer effects in humans. For example, in a key study recently conducted in humans, it was shown that natural killer cells with a highly mature effector phenotype are preferentially redistributed after exercise, and have the capacity to exert augmented cytotoxicity against myeloma and lymphoma cells *in vitro* (107, 108). In light of these results, research is now being conducted to harness the beneficial impact of acute exercise on lymphocyte kinetics for the purposes of cancer immunotherapy (121). It is beyond the scope of this review to discuss other findings in this exciting field and we briefly conclude that the aforementioned studies imply that exercise-induced lymphocytosis, and the lymphopenia that follows, is beneficial to the immune system’s capacity to identify and neutralize damaged and neoplastic cells in peripheral tissues. Furthermore, in the context of neoplastic growth, this process may be directly responsible for reduced incidence of cancer among physically active people across the lifespan (122). Further comprehensive discussion of the role of exercise and lymphocyte kinetics in anti-tumor responses can be found elsewhere (117). Clearly more research is needed in this area, and a shift in focus toward investigating the benefits—rather than purported detrimental effects—of exercise on health, is no doubt underway and will be a key focus for exercise immunologists in the coming years.

Transient Changes to Blood Lymphocyte Function in the Hours Following Exercise

A common misinterpretation, brought about by the exercise-induced reductions to blood lymphocyte frequency in the hours following exercise, is the observation that the functional capacity of immune cells in the peripheral blood is reduced in the hours following vigorous exercise. As measurements of cell function in peripheral blood are entirely dependent on the cells present at the time of sampling, a change to the composition of the cells in blood in the hours after exercise—as outlined in the section; “Transient Changes to Blood Lymphocyte Frequency in the Hours Following Exercise”—will consequently lead to parallel

changes in overall cell function, indicated by the performance of the sampled cells being assayed. For example, among CD8⁺ T cells, subsets that exhibit strong effector function (e.g., CD45 RA⁺CD27⁻CD28⁻CCR7⁻CD62L⁻CD57⁺), substantially increase during exercise (81, 123). Thus, during exercise, blood is predominantly occupied by cells capable of responding strongly (i.e., IFN-gamma production) to *in vitro* stimuli, and therefore, many studies have shown an increase in IFN-gamma production by cells isolated close to the exercise stimulus. In the hours following exercise, the same effector CD8⁺ T cells are subsequently redeployed to peripheral tissues, and, as such, this results in the blood having fewer cells capable of responding strongly to *in vitro* stimuli, thus explaining the commonly reported decrease in cellular function post-exercise. These effects have been neatly demonstrated approximately two decades ago when it was shown that IFN-gamma production by stimulated CD8⁺ T cells is reduced 2 h after completing a prolonged 2.5 h bout of cycling (124). Importantly, it was shown that this reduced capacity to produce IFN-gamma was due to a reduced number of IFN-gamma-positive CD8⁺ T cells in peripheral blood at the same time-point (124, 125).

The same principles apply to other cell functions, such as *in vitro* proliferation in response to mitogenic stimuli. However, with this measurement in particular, the commonly reported increase in T cell proliferation immediately after acute bouts of exercise is also strongly influenced by laboratory processes and *in vitro* assay conditions (e.g., blood processing, temperature, mitogen selection) (126). A recent meta-analysis of 24 studies concluded that lymphocyte proliferation is suppressed following acute bouts of exercise, and that a greater magnitude of suppression is caused by exercise lasting longer than 1 h, regardless of exercise intensity (127). However, this meta-analysis did not examine the most important determinant of cell function following exercise: the time-dependent changes in the cellular composition of the samples assayed. Thus, findings such as these should be interpreted with caution if analyses did not differentiate between studies collecting samples immediately after exercise or in the hours following.

As with research focusing on T cells, a similar group of studies citing reductions to natural killer cell cytotoxicity following acute exercise, reviewed elsewhere (128), did not always take into account dramatic shifts in the constitutional makeup of the natural killer cell compartment, which is known to change in response to exercise (81). Once more, changes to the functional capacity of the total natural killer cell pool are likely to have been misrepresented, given that many of these cells, with potent effector functions, are redistributed to peripheral tissues following exercise cessation. The principles discussed herein regarding lymphocyte function are also broadly applicable to the assessment of function in other cells, such as neutrophils and monocytes; the response of these cells to exercise is beyond the scope of this article and is reviewed elsewhere (8).

Taken together, it is important to emphasize that statements such as “acute intensive exercise elicits a depression of several aspects of acquired immune function” and “prolonged bouts of heavy exertion reduce the normal functioning of all major immune cell subtypes” mentioned elsewhere (8, 15) should be interpreted

with caution. We conclude that the results of studies exploring the effects of acute exercise on cell function must be considered very carefully in light of the time-dependent changes in the cellular composition of blood that typically arise following a vigorous bout.

Summary: Exercise Induces Lymphocyte Immune Surveillance Not Immune Suppression

In summary, strong evidence implies that a reduction in the frequency and function of lymphocytes (and other immune cells) in peripheral blood in the hours following vigorous and prolonged exercise does not reflect immune suppression. Instead, the observed lymphopenia represents a heightened state of immune surveillance and immune regulation driven by a preferential mobilization of cells to peripheral tissues. As such, nutritional interventions, which have been employed to dampen the magnitude of exercise lymphopenia (124, 129) are unlikely to reduce the incidence of infections, but interventions that augment exercise-induced lymphocyte trafficking may provide benefits (130).

PART B: REGULAR PHYSICAL ACTIVITY AND FREQUENT EXERCISE AUGMENT ASPECTS OF IMMUNE COMPETENCY ACROSS THE LIFESPAN

Contrary to a commonly held belief—outlined in Part A “Is it Time to Close the Shutters on the “Open Window” Hypothesis? A Bout of Exercise Does Not Suppress Immune Competency?”—that acute vigorous exercise can suppress aspects of immune function, an increasingly large body of research indicates that both single bouts of exercise, or frequent participation in regular exercise, can act as an adjuvant to stimulate the immune system. Numerous methods exist to assess the effects of behavioral interventions on immunity (131) but arguably the optimal means of evaluating global immune competency at a systems level is *via* assessment of the immune response to *in vivo* challenge, ideally with a novel and clinically recognized pathogen, for example *via* vaccination (132). Thus, here, in the first section of Part B “Regular Physical Activity and Frequent Exercise Augment Aspects of Immune Competency Across the Lifespan,” we focus on the influence that exercise has on immune responses to antigenic challenge, which requires a coordinated response from most, if not all, innate and adaptive immune system components. In light of the age-associated decline in immune competence with aging—caused in part by underlying changes to the numerical, phenotypic, and functional capacity of almost all innate and adaptive immune cells—the second section of Part B “Regular Physical Activity and Frequent Exercise Augment Aspects of Immune Competency Across the Lifespan,” will evaluate the impact aging has on the immune benefits that can be attained from exercise throughout the lifespan. A detailed discussion of immunological aging processes and the influence that an active lifestyle has on established biomarkers of an aging immune system is covered in Part C “Does exercise and Regular Physical Activity Influence Immunological Ageing?” of this article.

Exercise and Immune Responses to Experimental *In Vivo* Challenge Across the Lifespan

In a research context, the most clinically relevant model to assess *in vivo* challenge in a controlled manner is *via* vaccination. Vaccine administration assesses the integrated capacity of the immune system to recognize and process antigen, leading to antigen neutralization. In a clinical research context, vaccination responses are principally quantified clinically in two ways, either *via* antibody production from antigen-specific plasma cells or *via* cytokine responses—typically IFN-gamma production—from T cells stimulated with cognate antigen.

Vaccination: Effects of a Single Exercise Bout

Evidence from an array of studies, evaluated recently in a comprehensive review elsewhere (9), indicates that a single acute bout of exercise appears to enhance immune responses to vaccination in both younger and older individuals. The majority of studies to date have examined muscle-damaging upper arm resistance exercise performed close to the time of vaccination which is administered shortly after the regimen into exercised muscle. However, other modes of exercise, including acute bouts of whole body aerobic activity, have also been investigated. Six of the eight trials identified in the aforementioned review indicated a statistically significant exercise-induced enhancement of immune responses against constituent antigens contained within the vaccine administered (133–138). It is notable that in five of these studies, statistically significant benefits were found where the vaccine strains appeared to have lower immunogenicity (9). For example, it was shown in a trial of 133 young adults, approximately 20 years of age, receiving either a full- or half-dose Pneumovax-23 (a pneumococcal vaccine), that those who exercised at the time of receiving the half-dose vaccine had heightened responses to five of the eleven pneumococcal strains contained in the vaccine, whereas no differences were observed for the other six strains; and no benefits of exercise were observed for the full-dose vaccine (137). As such, given the potential effectiveness of exercise as an adjuvant in situations where vaccine immunogenicity is low, studying the effects of exercise on antibody responses in older adults—whom typically exhibit impaired responses—has received considerable attention. In one recent study, it was found that antibody responses in women 55–75 years of age, were significantly improved when moderate-intensity aerobic exercise was performed immediately prior to vaccination; however, no beneficial effects were found in men (138). Another trial reported no benefits of a single 45-min bout of moderate-intensity walking exercise on the immune response to influenza and pneumococcal vaccination in women around 47 years of age (139). Finally, a very recent study found no effect of a bout of resistance exercise on antibody responses to influenza vaccination in adults approximately 73 years of age (140). It is possible that a number of factors including immunological aging, biological sex and variations in sex hormone levels, and perhaps latent infections (e.g., herpes viruses) (141–143) limit the immunostimulatory effects of exercise, and many studies have not adequately controlled for these factors. Alternatively, in light of the mechanisms

proposed in the acute stress/exercise immune-enhancement hypothesis (93), it is plausible that more intensive exercise may be needed to elicit enhanced vaccine responses. Despite null findings, it is important to point out that few, if any, studies investigating the effects of acute exercise on vaccination responses have reported exercise-induced *impairment* to immune responses, and rather, these studies report that exercise has no effect, or in some cases a beneficial effect, on the immune response to vaccination in older adults.

Vaccination: Effects of Frequent Exercise Bouts

Data from vaccine studies exploring the effects of regular physical activity or frequent exercise training on the immune response to vaccination provides robust support for the argument that exercise enhances, rather than suppresses immunity. Indeed, at least eight studies have demonstrated heightened vaccination responses in older adults, typically over 60 years of age, who are physically active (144–151). For example, an early study categorized adults aged 62 years or older, into one of three groups: active (undertaking at least 20 min of vigorous exercise three or more times per week), moderately active (undertaking regular exercise but with lower intensity, frequency, and/or duration), or sedentary (non-exercisers). Two weeks after influenza vaccination, it was shown that serum anti-influenza IgG and IgM titers were higher in active versus sedentary adults, and so too were peripheral blood mononuclear cell responses to antigen-specific stimulation (144). In addition, a recent study has shown that men aged 65–85 years, who regularly undertook moderate or vigorous exercise training, exhibited higher antibody responses compared to inactive controls in response to an influenza vaccine (151). Data linking habitual levels of physical activity to enhanced immune competency in humans are supported by evidence from animal studies, and show that the immunological benefits of exercise may be particularly beneficial in enhancing otherwise poor responses in older age (152).

Interpreting Data From Vaccine Trials

A major criticism of vaccine and exercise trials conducted in humans is that many solely focus on the maximal antibody titer following vaccination, and it is not practical to follow-up with investigations into infection incidence as a gauge of protection status following vaccination (153). As such, it is unknown whether differences observed with absolute antibody titers, or the amount of IFN-gamma produced from stimulated T cells, between exercise and control groups, is representative of clinically meaningful benefits in terms of protection from infections. Three elegant studies in rodents imply that benefits, beyond antibody titer, may be brought about by acute exercise. In one of these studies, it was found that antibody responses to influenza exposure were *lower* in rodents that exercised around the time of exposure, compared to those that did not exercise, yet, exercised mice were still protected and did not exhibit any signs of infection upon re-exposure to the virus (154). Moreover, in an earlier study by the same authors, it was found that mice undertaking an acute bout of exercise before being exposed to influenza exhibited a lower severity of infection and had enhanced viral clearance and lower inflammation in the lungs in the days following

(155). Thus, it may be the case that exercise enhances immune responses, beyond those captured using maximal antibody titer as an endpoint. In accordance with this view, an elegant rodent study conducted by an independent group (156) found that mice exercised for 20–30 min at moderate intensity 4 h after intranasal influenza exposure had a substantially higher survival rate (18 of 22 survived; 82%) when compared to mice that did not exercise after influenza exposure (10 of 23 survived; 43%).

Contact Sensitivity Reactions and Acute Exercise Bouts

More recent studies have examined the effects of acute exercise on immune competency using other *in vivo* models of immune challenge that, in principle, also assess the coordinated efforts of immune system components. These studies have employed contact-sensitivity reactions by topically applying to skin, the dendritic cell and T cell stimulant (or attractant) diphenylcyclopropenone (DPCP), and the non-specific inflammatory stimulant, croton oil (157, 158). For example, in studies of young adults (approximately 20–30 years of age), by applying a primary sensitizing dose of DPCP 20 min after 2 h of moderate-intensity treadmill running, and assessing recall challenge 4 weeks later, it has been concluded that this form of exercise impairs both the induction of T cell immunity and the memory response (159, 160). Thirty minutes of moderate- or vigorous-intensity running had no effect, and no forms of exercise modulated the non-specific inflammatory challenge in response to croton oil (159, 160). Although these findings are biologically interesting, the clinical relevance of exercise-induced change is unclear, in part, because the process of DPCP-induced immune modulation is not well defined (158) unlike the immune response to antigen administration by vaccination.

Summary: Exercise Enhances Immune Responses to *In Vivo* Antigenic Challenge

We conclude that there is growing evidence from a powerful array of studies in humans and rodents, indicating that exercise enhances, or at least does not suppress immune responses to *in vivo* challenge in younger and older individuals. These observations—which contradict those predicated by the “open-window” hypothesis—support the contention that an acute bout of exercise has no detrimental immune consequences for health. Thus, exercise should be encouraged, particularly for older adults who are at greatest risk of infections and who may obtain the greatest exercise-induced benefits to immune competency; an overview of the impact of aging on the immunological benefits that can be attained from exercise is described next.

Does Aging Influence the Immunological Benefits of Exercise and Regular Physical Activity?

Effect of a Single Exercise Bout

Research investigating the effects of exercise on immune function has sought to establish whether the observed benefits, as outlined earlier in young adults, such as exercise-induced immune cell mobilisation, that has been implicated in protection against cancer, is also applicable to older adults. For example, it has been reported that the magnitude of T cell mobilization

in response to acute vigorous exercise is smaller in older (65 ± 1 years) compared to younger (22 ± 1 years) adults (161, 162). However, in this study, it was also shown that following exercise, the magnitude of T cell proliferation in response to mitogens was smaller in young adults, whereas a similar exercise-induced stimulation of natural killer cell cytotoxicity was observed for both groups (161, 162). It is beyond the scope of this review to fully critique investigations examining the influence of single exercise bouts on the function of different immune cells, with comparisons made between younger and older individuals across the lifespan; we refer the reader to comprehensive reviews on this topic (163–165). Nevertheless, it is important to point out that many studies in this area are difficult to interpret: at the time of their publication, the influence of *Cytomegalovirus* infection on the magnitude of exercise-induced immune responses was not known and was therefore not considered (123, 166). Moreover, while the magnitude of change to lymphocyte kinetics is likely to be important for detecting and eliminating viral and bacterial infections and neoplastic cells, this process is complex to study, and comparisons between younger and older people is difficult, partly due to other age-associated changes that influence the physiological response to exercise, such as the decline in fitness (e.g., sarcopenia, cardiorespiratory fitness) with age. It is likely that some, or all of these factors impact upon the efficacy of exercise as an adjuvant to vaccination responses in older adults, outlined earlier in the section “Exercise and Immune Responses to Experimental *In Vivo* Challenge Across the Lifespan.” In light of the challenge interpreting the clinical relevance of the aforementioned lymphocyte kinetics studies, from here onward, we briefly evaluate the impact of regular physical activity or frequent exercise bouts on immune competency across the lifespan, using measurements in samples collected from participants at rest.

Frequent Bouts of Exercise

Immune competency at rest has been assessed in cross-sectional studies, comparing elderly individuals differentiated by physical activity level or cardiorespiratory fitness, or by examining immune function before and after structured exercise training interventions. For example, it has commonly been reported among the elderly, that the most active participants, compared to those who are least active, show the highest T cell proliferation and cytokine production in response to mitogens (163–165). Fewer studies have assessed innate immune competency, but higher natural killer cell cytotoxicity has been consistently shown among the elderly who are active compared to less active age-matched controls (163–165). Recent studies have expanded measurements into other innate immune cells such as neutrophils. For example, a recent cross-sectional study of 211 elderly adults, showed that neutrophils from the 20 most active participants, compared to the 20 least active participants, migrated toward interleukin (IL)-8 with greater chemotactic accuracy, but there were no differences in chemotactic speed (167). In addition, a recent exercise training study has shown in both young and middle-aged adults, that 10 weeks of moderate-intensity continuous cycling training, or high-intensity interval cycling training, improve neutrophil and monocyte phagocytosis and oxidative burst irrespective

of age (168). Improvements in these common measurements of immune competency, however, are not always consistent in longitudinal studies employing exercise training interventions, with around half of studies reporting improvements, and half reporting no change (163–165). One reason for this could be because the dramatic effects of *Cytomegalovirus* on driving immunological aging was not considered by most of these studies, and it is feasible that results would be different when examining individuals who are latently infected compared to those who are seronegative. Importantly however, no studies report impaired immune competency from increased participation in structured exercise. Altogether, we conclude that despite declines in fitness and immune competency, aging does not appear to negate the immunological benefits that can be attained from exercise, and indeed, frequent participation in exercise across the lifespan may lead to immune benefits, even in older age.

PART C: DOES EXERCISE AND REGULAR PHYSICAL ACTIVITY INFLUENCE IMMUNOLOGICAL AGEING?

Since the first exercise immunology research in the early 1900s, and the substantial increase in scientific interest from the late 1980s and early 1990s (169), studies examining interaction between immune function and lifestyle factors such as exercise and physical activity have become common. Although a few exercise studies published in the last 10–15 years have investigated some immunological processes relevant to aging, health, and disease, the theme of exercise-induced “immune suppression” continues to influence the design of new studies or at least features in the interpretation of findings and is often justified or contextualized with relevance to self-reported illness symptoms among athletes. As outlined in Part A “Is it Time to Close the Shutters on the “Open-Window” Hypothesis? A Bout of Exercise Does Not Suppress Immune Competency,” there is limited reliable evidence to show that exercise heightens risk of opportunistic infections, but there is, however, a growing body of evidence to show that exercise enhances, rather than suppresses, immune competency, as summarized in Part B: “Regular Physical Activity and Frequent Exercise Augment Aspects of Immune Competency Across the Lifespan.” The beneficial effects of exercise on immune function are likely to be greatest for elderly people exhibiting the age-associated deterioration of immune competency, also referred to as immunosenescence. Moreover, preliminary evidence suggests that physical activity and regular structured exercise may even limit or delay immunological aging. Here, in Part C: “Does Exercise and Regular Physical Activity Influence Immunological Ageing?” we evaluate whether an active lifestyle influences immunosenescence.

The Aging Immune System

Aging is associated with profound changes to the numerical, phenotypic, and functional capacity of almost all innate and adaptive immune cells, resulting in altered immune responses. Some innate immune cells exhibit numerical, phenotypic, and functional alterations with aging, whereas others appear to be less affected (170).

For example, the numbers and functions of eosinophils, basophils, and mast cells appear to be largely unchanged with aging, or at least, there is not a clear effect of age based on the limited current literature (170). Neutrophil numbers often increase with aging, but these cells exhibit diminished phagocytosis and impaired chemotaxis, although chemokinesis is maintained (170). Natural killer cells increase with aging, driven by an accumulation of cytotoxic CD56^{dim} cells but a decline in regulatory CD56^{bright} cells, and overall, cytokine production and cytotoxicity are less on a per cell basis (170). Other innate lymphocytes, such as invariant CD1d-restricted natural killer T cells (iNKT cells), which represent <1% of the T cell pool, decline in number but age-associated changes to their function have not been established (171). Monocyte numbers are stable with aging, but classical cells (CD14⁺⁺CD16⁻) decline, and intermediate (CD14⁺CD16⁺) and non-classical (CD14⁺CD16⁺⁺) cells increase, but overall, monocyte cytokine production is impaired (170, 172). These changes with blood monocytes are thought to be mirrored by tissue-resident macrophages, whereby classically activated M1 cells decline, and alternatively activated M2 cells accumulate (173). However, alterations in tissue-resident cells with advancing age are very likely to be a result of adipose tissue accumulation and dysfunction that also occurs in parallel with aging (174, 175). Indeed, inflamed adipose tissue attracts macrophages with cell-surface characteristics similar to M2 alternatively activated cells—often assumed to be anti-inflammatory. However, despite their cell-surface phenotype, these cells are potent producers of inflammatory cytokines in adipose tissue, and likely drive age- and obesity-associated inflammation (176–179). Thus, the M1/M2 paradigm for macrophages is likely to be an over-simplification (180, 181). It is unknown if other, primarily tissue-resident cells, are affected by adipose tissue dysfunction, but with aging, the number and function of dendritic cells have been reported to decrease in the skin and mucosal membranes (170). There is also an age-associated increase in myeloid-derived suppressor cells—a heterogeneous population of granulocytes, macrophages, and dendritic cells—that may impair aspects of immune function by producing reactive oxygen species and inhibitory cytokines (182).

Within the adaptive immune system, there are substantial changes to the numbers, function, and phenotype of T cells with aging. Among the broad population of CD4⁺ T-helper cells, aging is associated with a predominance of Th2 (i.e., IL-4 and IL-10), and Th17-producing cells (i.e., IL-17-producing cells that are associated with autoimmune disease), whereas there is a decline in cells with a Th1 profile [i.e., IFN-gamma- and tumor necrosis factor-alpha (TNF-alpha)-producing cells] (183, 184). With aging, the numbers and proportions of antigen-inexperienced CD4⁺ and CD8⁺ T cells decreases (e.g., CD27⁺CD28⁺CD45RA⁺CD57⁻CD62L⁺CCR7⁺KLRG1⁻ naïve cells) (185, 186). In parallel, the numbers and proportions of antigen-experienced CD4⁺ and CD8⁺ T cells increases (e.g., CD27⁻CD28⁻CD45RA⁺CD57⁺CD62L⁻CCR7⁻KLRG1⁺ memory cells), and these cells are potent producers of inflammatory cytokines (185, 186). These changes are driven by lower hematopoietic stem cell numbers, thymic involution resulting in reduced output of antigen-naïve T cells, and infection with latent viruses, in particular *Cytomegalovirus* (185, 187). With aging, T cells that express natural killer cell-associated cell-surface proteins (NKT-like cells) also accumulate,

exhibiting similar changes to their phenotype, functional properties, and specificities as with the broader population of CD4⁺ and CD8⁺ T cells (171). There is an age-associated decline in the total number of $\gamma\delta$ T cells; however age *per se*, in the absence of chronic infections, is associated with a decline in V δ 2 cells (60–80% of $\gamma\delta$ T cells), whereas V δ 1 (15–20% $\gamma\delta$ T cells) remain stable (188). Some evidence suggests that $\gamma\delta$ T cell proliferative responses are impaired with aging, perhaps due to increased susceptibility of V δ 2 cells to apoptosis (189, 190). Natural regulatory T cells increase with aging whereas inducible regulatory T cells decrease, but it is unclear if their function is affected (191). As with T cells, aging is associated with a decline in the numbers and proportions of naïve B cells, an accumulation of memory B cells with limited specificities, and impaired plasma cell antibody production (192).

Several robust and accepted hallmarks of immunosenescence have been established, especially within the adaptive immune system. For example, low numbers and proportions of naïve T cells (in particular CD8⁺ T cells) and high numbers and proportions of memory T cells (especially late-stage differentiated CD8⁺ T cells) are well established biomarkers (185, 186, 193, 194). In addition, a cluster of parameters, revealed in longitudinal studies of octogenarians and nonagenarians from an isolated population in Sweden, have been referred to as the immune risk profile (195–197). Biomarkers included low numbers and proportions of B cells, high numbers and proportions of late-stage differentiated CD8⁺ T cells (i.e., CD27⁻CD28⁻), poor T cell proliferation in response to mitogens, a CD4:CD8 ratio of <1.0, infection with *Cytomegalovirus* and high plasma IL-6, which together, predicted greater all-cause mortality at 2-, 4-, and 6-year follow-up (195, 197–200). Indeed, the age-associated increase in systemic inflammation, referred to as “inflammaging” is another principle observation among aging and longevity studies (201, 202). Subsequently, high levels of IL-6, TNF-alpha, and C-reactive protein, have been associated with shorter survival (203–205). Overall, it is well established that elderly individuals exhibit impaired immune responses to *in vivo* challenge with novel antigens (143, 206, 207) and these individuals are subsequently thought to be at increased risk of infection. Encouragingly however, as outlined in Part B: “Regular Physical Activity and Frequent Exercise Augment Aspects of Immune Competency Across the Lifespan” exercise can be a potent stimulus of immune function, including the response to vaccination, and some evidence suggests that exercise might delay or limit the age-associated decline in immune competency.

Relationships Between Exercise and Regular Physical Activity With Hallmarks of an Aging Adaptive Immune System

As summarized in Part B: “Regular Physical Activity and Frequent Exercise Augment Aspects of Immune Competency Across the Lifespan” many studies have examined the influence of regular physical activity or frequent structured exercise on the function of the adaptive immune system with aging (163–165). Here, we focus on recent studies that have examined relationships between exercise, physical activity or cardiorespiratory fitness, and the numbers and proportions of CD4⁺ and CD8⁺ naïve and memory

T cells, as hallmarks of immunosenescence. Indeed, a small number of studies have investigated whether the characteristics of the T cell pool are influenced by an active lifestyle. There is a larger body of evidence in young adults, typically between 18 and 30 years, compared to older adults, hereafter considered as being over 40 years of age due to the characteristics of studies published to date. If an active lifestyle can be linked with a smaller accumulation of memory T cells, and a smaller decline in naïve T cells, then in young adults, one interpretation might be that exercise prevents, or at least delays immunosenescence, whereas in older adults, these associations could be interpreted as countering or limiting the development of an age-associated immune profile.

Experimental Evidence in Young People: Can Exercise Prevent or Delay Aging of the Adaptive Immune System?

The characteristics of the T cell pool have been examined in 16 young adults (50% male; 18 ± 2 years) classified as being very active (well-trained soccer players self-reporting 9–12 h of exercise per week) and compared to 16 young adults (50% male; 19 ± 2 years) classified as being untrained (individuals self-reporting 2–3 h of exercise per week). Untrained individuals showed the highest proportions of CD4⁺ and CD8⁺ memory T cells, and the lowest proportions of CD8⁺ naïve T cells, defined on the basis of CD57 and CD28 expression (208). Although these results suggest that regular exercise might limit the age-associated accumulation of memory T cells and decline in naïve T cells, the effects were strongly influenced by sex: only untrained males exhibited high proportions of memory T cells and low proportions of naïve cells compared to trained males, and these effects were driven by changes in the CD4⁺ T cell pool (208). Extending these findings by examining a female-only population of young adults, the same authors compared 13 well-trained soccer players (self-reporting around 12 h of exercise per week; 20 ± 2 years) to 13 untrained controls (self-reporting around 3 h of exercise per week; 21 ± 2 years) (209). Trained females exhibited a greater proportion of CD8⁺ naïve T cells compared to untrained females, but these associations did not remain statistically significant after controlling for body fat percentage (trained 21.7 ± 4.0 versus untrained $25.1 \pm 4.1\%$, $p < 0.05$) (209). Indeed, very few studies have considered whether relationships between an active lifestyle and markers of an aging immune system could be influenced by other factors, such as body composition. Recently, a very small study of 15 males aged 30 ± 4 years, categorized participants using a combination of gold standard methods for measuring physiological and lifestyle variables (210). Three groups were formed (sedentary, active, and very active) on the basis of objectively assessed habitual physical activity, directly measured cardiorespiratory fitness, and body composition assessed with dual energy X-ray absorptiometry (210). This work showed that sedentary individuals had higher proportions of memory CD4⁺ T cells expressing CD45RO and PD-1, supporting the results of other published studies that have not taken body composition into consideration.

It should be emphasized, that among younger adults in particular, mixed results have been reported when investigating links between an active lifestyle and hallmarks of an aging

immune system. Most investigations have been cross-sectional in design, or have made observations between groups over short periods. For example, one study has shown that national standard triathletes (age range 18–36 years, $n = 19$), appear to exhibit impaired thymic output, assessed by T cell receptor excision circle levels (211). Among CD4⁺ and CD8⁺ T cells, these athletes had lower absolute numbers of naïve cells and higher absolute numbers of memory cells compared to age-matched less active controls ($n = 16$), as shown by CD45RA, CD45RO, and CD27 expression (211). Similar observations have been made by examining individuals in the third decade of life, showing that endurance athletes, compared to less active controls, appear to exhibit slightly larger accumulations of memory T cells and slightly fewer naïve T cells, defined by CD45RA and CCR7 expression (212). Thus, it appears that among younger individuals (i.e., less than 40 years of age), if exercise is undertaken at very extreme volumes, such as more than 5–10 times the amount of physical activity recommended each week (i.e., 12–25 h per week), this might contribute toward a small decline in naïve T cells and a small increase in memory T cells. It is possible that these changes are due to reactivation of latent viruses, which could be independent of immune function, and driven by exercise-induced adrenergic activity, oxidative stress and inflammatory cytokines (48–50). However, these results might also be explained by fluctuations in cell numbers and cell sub-populations in peripheral blood over time. Such changes have been interpreted as being linked to exercise training load (213, 214), but it is also conceivable that these changes occur due to seasonal variation, as has been shown in non-exercise contexts (215, 216).

Experimental Evidence in Older People: Can Exercise Limit or Counter Aging of the Adaptive Immune System?

Although most studies have examined associations between biomarkers of an aging adaptive immune system in young adults, other studies have made measurements across a broader range of ages. For example, one study examined 102 men ranging in age from 18 to 61 years (mean 39 ± 6 years) (217). It was shown that the proportion of the CD4⁺ and CD8⁺ T cell pool comprising of memory cells (defined as KLRG1⁺CD57⁺ or KLRG1⁺CD28⁻) was inversely correlated with cardiorespiratory fitness, which is largely indicative of an active lifestyle (217). The age-associated decrease of naïve T cells (defined as KLRG1⁻CD57⁻ or KLRG1⁻CD28⁺) and increase in memory T cells did not withstand statistical adjustment for cardiorespiratory fitness, but remained significant after adjusting statistically for body composition and *Cytomegalovirus* infection (217). Thus, it was concluded that fitter individuals exhibit a smaller age-associated decline of naïve T cells and a smaller accumulation of memory T cells.

As with work examining relationships between an active lifestyle and hallmarks of an aging adaptive immune system in young and middle-aged adults, similar associations have been shown in an older population of 61 men aged 65–85 years (218). In this study, participants self-reported to be “untrained” ($n = 15$), or to lead a “moderate” training lifestyle ($n = 16$;

taking part in team sports or running less than 6 km two to three times per week), or an “intense” training lifestyle ($n = 15$; running approximately 10 km at least 5 days per week). These categories were confirmed with a validated physical activity questionnaire and by measurement of cardiorespiratory fitness. Both training groups exhibited a lower proportion of CD4⁺ and CD8⁺ memory cells (defined as CD45RA⁺CCR7⁻), but these associations were largest and only statistically significant among men leading an “intense” training lifestyle, suggesting a dose–response effect of exercise. Although findings were less clear when examining other cell subpopulations based on CD45RA and CCR7 expression, and there were no effects of exercise when examining CD28⁻ cells, men in the “trained” groups had T cells with the longest telomeres (218). Another recent study compared 125 adults (55–79 years of age) who maintained a high level of cycling throughout life to 75 age-matched inactive controls (219). Within the CD4⁺ and CD8⁺ T cell pool, the frequency of naïve cells (defined as CD45RA⁺) was greater, and the frequency of memory cells (defined as CD45RA⁻) was lower among cyclists. Extended phenotyping revealed that CD4⁺ and CD8⁺ CCR7-CD45RA⁺ accumulation was less among cyclists, but no differences were found for CD28-CD57⁺ cells. Cyclists also exhibited higher frequencies of recent thymic emigrants and regulatory B cells, lower Th17 polarization, and in plasma, higher IL-7 and lower IL-6 (219). Despite these findings, suggesting a beneficial effect of leading an active lifestyle on immunosenescence among older adults, there is some inconsistency in the literature. For example, comparing elderly athletes ($n = 12$, approximately 74 years of age) to less active age-matched controls ($n = 26$), there were no differences in thymic output, the proportions of naïve or memory CD4⁺ and CD8⁺ T cells (defined with CD45RA and CCR7 expression), or T cell activation in response to anti-CD3 stimulation (212). Most investigations of T cell immunosenescence and lifestyle among healthy elderly adults have had cross-sectional study designs. Longitudinal studies, or randomized and controlled trials of exercise training are lacking and might yield promising results. For example, one study has compared 6 months of exercise training in men and women ($n = 28$, aged 61–76 years) to a similar group who maintained their current lifestyle ($n = 20$, aged 62–79 years) (220). Using a simple immunophenotyping strategy, the results showed that the proportion of CD4⁺ T cells expressing CD28 increased in the exercise group after 6 months, but not in those who maintained their lifestyle (220).

Summary of Experimental Evidence

In summary, evidence shows that the characteristics of the T cell pool appear to be influenced by leading an active lifestyle, determined by exercise training, physical activity level, or cardiorespiratory fitness. It seems that among both the young and elderly, an active lifestyle is generally linked to lower numbers and proportions of memory T cells and higher numbers and proportions of naïve T cells (10). This summary is partly supported by a recent systematic review, concluding that regular structured exercise increases the number of naïve T cells in peripheral blood at rest (221). Altogether, findings

from recent studies examining relationships between an active lifestyle and the characteristics of the T cell pool—as robust and accepted biomarkers of immunosenescence—support observations from some cross-sectional and longitudinal studies, showing that other measures of immune competency, which typically decline with aging, can be improved with physical activity or regular structured exercise (163–165). However, further research is needed in this area that employs precise lifestyle measurements (e.g., using wearable technology to assess physical activity, and dual energy X-ray absorptiometry to measure body composition) and more robust measurements of immune competency (e.g., absolute cell counts rather than proportions, measurements of cell function, and *in vivo* antigen challenges) while controlling for factors that drive immunosenescence (e.g., inflammation and *Cytomegalovirus* infection).

Mechanisms

Links between a physically active lifestyle with lower numbers or proportions of memory T cells, and higher numbers or proportions of naïve T cells, have been hypothesized as being driven by the acute effects of exercise bouts. For example, it has been suggested that repeated bouts of exercise might prevent or delay immunological aging by limiting the accumulation of CD4⁺ and CD8⁺ antigen-experienced memory T cell clones, repopulating blood with antigen-inexperienced naïve T cells (11, 12). In this hypothesis, it is proposed that memory T cells are frequently mobilized into blood during regular bouts of exercise, followed by an extravasation to peripheral tissues, where these cells are exposed to pro-apoptotic stimuli, such as reactive oxygen species, glucocorticoids, and cytokines (11, 12). Subsequently, it is proposed that the number of naïve T cells increases as part of a negative feedback loop governing the number of naïve and memory cells in the immune system, which is bolstered by exercise-induced thymopoiesis and extra-thymic T cell development (11, 12). Supporting the mechanisms proposed in this hypothesis, many investigations have shown that memory T cells are mobilized into the circulation during exercise, followed by extravasation out of the bloodstream in the hours following (81, 123). In addition, studies in mice show that lymphocyte apoptosis occurs post-exercise in tissues thought to be the homing destination of mobilized cells (222). Although some T cells mobilized by exercise might be resistant to apoptosis, given that *Cytomegalovirus*-specific CD8⁺ T cells express high levels of Bcl-2 (223), other work has shown that *Cytomegalovirus*-specific CD8⁺ T cells, are equally as susceptible to Fas-induced apoptosis as the total pool of CD8⁺ T cells (224). Further, irrespective of virus specificity, studies have shown that T cells expressing cell-surface proteins such as CD57 and KLRG1 are more susceptible to H₂O₂-induced apoptosis than total lymphocytes and naïve T cells (225, 226). Thus, the concept of exercise directly countering memory T cell accumulation is supported by evidence from human and animal studies.

It is unknown whether triggering apoptosis among expanded clones of memory T cells specific for viruses such as *Cytomegalovirus* is advantageous. For example, in a transplant

setting, *Cytomegalovirus* disease occurs when T cells fail to provide antiviral control (227) and a robust pro-inflammatory response to *Cytomegalovirus* has been associated with longer survival in the elderly (228). However, it remains to be determined what proportion of the T cell pool needs to be specific for *Cytomegalovirus* to limit viral reactivation. Infection with *Cytomegalovirus* results in approximately 10% of the CD4⁺ and CD8⁺ T cell pool becoming specific for this virus (229), although large inter-individual differences exist. For example, it has been reported that 23% of the CD8⁺ T cell pool can become specific for a single *Cytomegalovirus* epitope (223). Traditionally, it has been considered disadvantageous for such a large proportion of the T cell pool to be specific for one virus. This view is linked to another age- or infection-associated change that occurs in parallel—a fall in the numbers and proportions of naïve cells—which has been interpreted as limiting capacity to engage novel antigens. These interpretations are based upon two assumptions. First, there is an upper limit to the size of the immune system, and second, thymic output is negligible after adolescence (230). Thus, it has been proposed that antigen-inexperienced naïve T cells could be “used up” due to ongoing differentiation into antigen-experienced memory T cells that “fill up” immunological “space” (230). It has also been proposed that this accumulation of antigen-experienced memory T cells leads to “squeezing out” of T cells targeting less dominant viruses leading to loss of viral control (231). This concept of a fixed amount of immunological space has since been debated (232, 233) and thymic output is now known to persist, albeit reduced, up until around 70 years of age (234). However, even if removal of some memory T cells is not essential for maintaining an effective T cell pool, assuming these cells contribute to systemic inflammation, their removal might limit “inflammaging” (230). Despite uncertainties over the susceptibility of memory T cells to undergo apoptosis, or whether it is advantageous to stimulate their removal, it seems that exercise-induced immune cell death in the tissues has relevance to other processes. For example, apoptotic cells and cell debris stimulates hematopoietic stem cell mobilization into blood (95) perhaps promoting trafficking to the thymus or extrathymic sites facilitating output of naïve T cells (235). Additional support for exercise stimulating production of naïve T cells is provided by work showing that contracting skeletal muscle produces IL-7 (236) which might increase thymic mass and function (237).

A physically active lifestyle might also counter T cell immunosenescence indirectly, perhaps by limiting adipose tissue accumulation and dysfunction that occurs with aging and obesity (174, 238, 239). Indeed, obesity has been linked with impaired lymphocyte proliferation (240), shorter leukocyte telomere length (241), and a skewing of the T cell pool toward a regulatory and Th2-phenotype (242). In addition, large expansions of differentiated $\alpha\beta$ T cells and $\gamma\delta$ T cells have been shown among people with obesity, with $\gamma\delta$ T cells exhibiting impaired antiviral function (243–245). It is generally accepted that repeated stimulation with antigens from *Cytomegalovirus* drives immunosenescence (185, 186, 193, 194). With obesity, adipose tissue is the primary source of pro-inflammatory cytokines and reactive oxygen species (174, 246) which can reactivate *Cytomegalovirus* directly (48, 49).

Thus, exercise might limit T cell immunosenescence by decreasing visceral and subcutaneous adipose tissue (238), providing a potent anti-inflammatory and anti-oxidative stimulus (247, 248). In turn, lower systemic inflammation and better redox balance might limit viral reactivation, reducing stimulation with antigens from viruses such as *Cytomegalovirus*. In addition, T cell dysfunction might also be prevented, in part, by limiting reactive oxygen species production (249).

In summary, leading a physically active lifestyle appears to limit the age-associated changes to the cellular composition of the adaptive immune system, but the mechanisms are yet to be determined. Exercise might counter the expansion of memory T cells directly, which is desirable assuming these cells contribute to systemic inflammation and not all are required to control latent viruses. Limiting the expansion of memory T cells also assumes the “size” of the immune system is fixed, the capacity to produce antigen-naïve T cells is limited, and these constraints contribute to immune decline in the elderly. However, exercise might affect memory T cell accumulation indirectly, by reducing viral reactivation, or preventing T cell senescence, by controlling adipose tissue deposition and dysfunction that drives inflammation and oxidative stress with aging and obesity.

CONCLUDING REMARKS

Contemporary evidence from epidemiological studies shows that leading a physically active lifestyle reduces the incidence of communicable (e.g., bacterial and viral infections) and non-communicable diseases (e.g., cancer), implying that immune competency is enhanced by regular exercise bouts. However, to this day, research practice, academic teaching, and even physical activity promotion and prescription continues to consider a prevailing myth that exercise can temporarily suppress immune function. We have critically reviewed related evidence, and conclude that regular physical activity and frequent exercise are beneficial, or at the very least, are not detrimental to immunological health. We summarize that (i) limited reliable evidence exists to support the claim that exercise suppresses cellular or soluble immune competency, (ii) exercise *per se* does not heighten the risk of opportunistic infections, and (iii) exercise can enhance *in vivo* immune responses to bacterial, viral, and other antigens. In addition, we present evidence showing that regular physical activity and frequent exercise might limit or delay immunological aging. We conclude that leading an active lifestyle is likely to be beneficial, rather than detrimental, to immune function, which may have implications for health and disease in older age.

AUTHOR CONTRIBUTIONS

JC and JT contributed equally toward literature searching and retrieval, the ideas and interpretation of the studies described, drafting and revision of the manuscript, and approval of the final version to be published. JT and JC both agreed to be accountable for the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

REFERENCES

- Warburton DER, Bredin SSD. Health benefits of physical activity: a systematic review of current systematic reviews. *Curr Opin Cardiol* (2017) 32(5): 541–56. doi:10.1097/HCO.0000000000000437
- Pape K, Ryttergaard L, Rotevatn TA, Nielsen BJ, Torp-Pedersen C, Overgaard C, et al. Leisure-time physical activity and the risk of suspected bacterial infections. *Med Sci Sports Exerc* (2016) 48(9):1737–44. doi:10.1249/MSS.0000000000000953
- Kostka T, Berthouze SE, Lacour J, Bonnefoy M. The symptomatology of upper respiratory tract infections and exercise in elderly people. *Med Sci Sports Exerc* (2000) 32(1):46–51. doi:10.1097/00005768-200001000-00008
- Leveille SG, Gray S, LaCroix AZ, Ferrucci L, Black DJ, Guralnik JM. Physical inactivity and smoking increase risk for serious infections in older women. *J Am Geriatr Soc* (2000) 48(12):1582–8. doi:10.1111/j.1532-5415.2000.tb03867.x
- Romaniszyn D, Pobiega M, Wojkowska-Mach J, Chmielarczyk A, Gryglewska B, Adamski P, et al. The general status of patients and limited physical activity as risk factors of methicillin-resistant *Staphylococcus aureus* occurrence in long-term care facilities residents in Krakow, Poland. *BMC Infect Dis* (2014) 14:271. doi:10.1186/1471-2334-14-271
- Baik I, Curhan GC, Rimm EB, Bendich A, Willett WC, Fawzi WW. A prospective study of age and lifestyle factors in relation to community-acquired pneumonia in US men and women. *Arch Intern Med* (2000) 160(20):3082–8. doi:10.1001/archinte.160.20.3082
- Walsh NP, Gleeson M, Pyne DB, Nieman DC, Dhabhar FS, Shephard RJ, et al. Position statement. Part two: maintaining immune health. *Exerc Immunol Rev* (2011) 17:64–103.
- Walsh NP, Gleeson M, Shephard RJ, Gleeson M, Woods JA, Bishop NC, et al. Position statement. Part one: immune function and exercise. *Exerc Immunol Rev* (2011) 17:6–63.
- Pascoe AR, Fiatarone Singh MA, Edwards KM. The effects of exercise on vaccination responses: a review of chronic and acute exercise interventions in humans. *Brain Behav Immun* (2014) 39:33–41. doi:10.1016/j.bbi.2013.10.003
- Turner JE. Is immunosenescence influenced by our lifetime “dose” of exercise? *Biogerontology* (2016) 17(3):581–602. doi:10.1007/s10522-016-9642-z
- Simpson RJ, Guy K. Coupling aging immunity with a sedentary lifestyle: has the damage already been done? – a mini-review. *Gerontology* (2010) 56(5):449–58. doi:10.1159/000270905
- Simpson RJ. Aging, persistent viral infections, and immunosenescence: can exercise “make space”? *Exerc Sport Sci Rev* (2011) 39(1):23–33. doi:10.1097/JES.0b013e318201f39d
- Turner JE, Brum PC. Does regular exercise counter T cell immunosenescence reducing the risk of developing cancer and promoting successful treatment of malignancies? *Oxid Med Cell Longev* (2017) 2017:4234765. doi:10.1155/2017/4234765
- Mackinnon LT. *Advances in Exercise Immunology*. Champaign, IL: Human Kinetics (1999). xii, 363 p.
- Peake JM, Neubauer O, Walsh NP, Simpson RJ. Recovery of the immune system after exercise. *J Appl Physiol* (1985) (2017) 122(5):1077–87. doi:10.1152/jappphysiol.00622.2016
- Cowles WN. Fatigue as a contributory cause of pneumonias. *Boston Med Surg J* (1918) 179:555. doi:10.1056/NEJM191810311791801
- Peters EM, Bateman ED. Ultramarathon running and upper respiratory tract infections. An epidemiological survey. *S Afr Med J* (1983) 64(15):582–4.
- Nieman DC, Johanssen LM, Lee JW, Arabatzis K. Infectious episodes in runners before and after the Los Angeles Marathon. *J Sports Med Phys Fitness* (1990) 30(3):316–28.
- Nieman DC, Johanssen LM, Lee JW. Infectious episodes in runners before and after a roadrace. *J Sports Med Phys Fitness* (1989) 29(3):289–96.
- Spence L, Brown WJ, Pyne DB, Nissen MD, Sloots TP, McCormack JG, et al. Incidence, etiology, and symptomatology of upper respiratory illness in elite athletes. *Med Sci Sports Exerc* (2007) 39(4):577–86. doi:10.1249/mss.0b013e31802e851a
- Gleeson M. Immune function in sport and exercise. *J Appl Physiol* (1985) (2007) 103(2):693–9. doi:10.1152/jappphysiol.00008.2007
- Cox AJ, Gleeson M, Pyne DB, Callister R, Hopkins WG, Fricker PA. Clinical and laboratory evaluation of upper respiratory symptoms in elite athletes. *Clin J Sport Med* (2008) 18(5):438–45. doi:10.1097/JSM.0b013e318181e501
- Makela MJ, Puhakka T, Ruuskanen O, Leinonen M, Saikku P, Kimpimaki M, et al. Viruses and bacteria in the etiology of the common cold. *J Clin Microbiol* (1998) 36(2):539–42.
- Nieman DC. Exercise, infection, and immunity. *Int J Sports Med* (1994) 15(Suppl 3):S131–41. doi:10.1055/s-2007-1021128
- Meeusen R, Duclos M, Foster C, Fry A, Gleeson M, Nieman D, et al. Prevention, diagnosis and treatment of the overtraining syndrome: joint consensus statement of the European College of Sport Science (ECSS) and the American College of Sports Medicine (ACSM). *Eur J Sport Sci* (2013) 13(1):1–24. doi:10.1080/17461391.2012.730061
- Meeusen R, Duclos M, Foster C, Fry A, Gleeson M, Nieman D, et al. Prevention, diagnosis, and treatment of the overtraining syndrome: joint consensus statement of the European College of Sport Science and the American College of Sports Medicine. *Med Sci Sports Exerc* (2013) 45(1):186–205. doi:10.1249/MSS.0b013e318279a10a
- Edwards JR, Walsh NP, Diment PC, Roberts R. Anxiety and perceived psychological stress play an important role in the immune response after exercise. *Exerc Immunol Rev* (2018) 24:26–34.
- Glaser R, Kiecolt-Glaser JK. Stress-induced immune dysfunction: implications for health. *Nat Rev Immunol* (2005) 5(3):243–51. doi:10.1038/nri1571
- Kohut ML, Lee W, Martin A, Arnston B, Russell DW, Ekkekakis P, et al. The exercise-induced enhancement of influenza immunity is mediated in part by improvements in psychosocial factors in older adults. *Brain Behav Immun* (2005) 19(4):357–66. doi:10.1016/j.bbi.2004.12.002
- Bermon S, Castell LM, Calder PC, Bishop NC, Blomstrand E, Mooren FC, et al. Consensus statement immunonutrition and exercise. *Exerc Immunol Rev* (2017) 23:8–50.
- Choudhry AJ, Al-Mudaimigh KS, Turkistani AM, Al-Hamdan NA. Hajj-associated acute respiratory infection among hajjis from Riyadh. *East Mediterr Health J* (2006) 12(3–4):300–9.
- Svensden IS, Taylor IM, Tonnessen E, Bahr R, Gleeson M. Training-related and competition-related risk factors for respiratory tract and gastrointestinal infections in elite cross-country skiers. *Br J Sports Med* (2016) 50(13):809–15. doi:10.1136/bjsports-2015-095398
- Gleeson M. Immunological aspects of sport nutrition. *Immunol Cell Biol* (2016) 94(2):117–23. doi:10.1038/icb.2015.109
- Schwellnus MP, Derman WE, Jordaan E, Page T, Lambert MI, Readhead C, et al. Elite athletes travelling to international destinations >5 time zone differences from their home country have a 2–3-fold increased risk of illness. *Br J Sports Med* (2012) 46(11):816–21. doi:10.1136/bjsports-2012-091395
- Fondell E, Lagerros YT, Sundberg CJ, Lekander M, Balter O, Rothman KJ, et al. Physical activity, stress, and self-reported upper respiratory tract infection. *Med Sci Sports Exerc* (2011) 43(2):272–9. doi:10.1249/MSS.0b013e3181edf108
- Martensson S, Nordebo K, Malm C. High training volumes are associated with a low number of self-reported sick days in elite endurance athletes. *J Sports Sci Med* (2014) 13(4):929–33.
- Hellard P, Avalos M, Guimaraes F, Toussaint JF, Pyne DB. Training-related risk of common illnesses in elite swimmers over a 4-yr period. *Med Sci Sports Exerc* (2015) 47(4):698–707. doi:10.1249/MSS.0000000000000461
- Hoffman MD, Krishnan E. Health and exercise-related medical issues among 1,212 ultramarathon runners: baseline findings from the Ultrarunners Longitudinal TRacking (ULTRA) study. *PLoS One* (2014) 9(1):e83867. doi:10.1371/journal.pone.0083867
- Hoffman MD, Fogard K. Demographic characteristics of 161-km ultramarathon runners. *Res Sports Med* (2012) 20(1):59–69. doi:10.1080/15438627.2012.634707
- Malm C. Susceptibility to infections in elite athletes: the S-curve. *Scand J Med Sci Sports* (2006) 16(1):4–6. doi:10.1111/j.1600-0838.2005.00499.x
- Derman W, Schwellnus M, Jordaan E. Clinical characteristics of 385 illnesses of athletes with impairment reported on the WEB-IISS system during the London 2012 Paralympic Games. *PM R* (2014) 6(8 Suppl):S23–30. doi:10.1016/j.pmrj.2014.05.016
- Bonini M, Gramiccioni C, Fioretti D, Ruckert B, Rinaldi M, Akdis C, et al. Asthma, allergy and the Olympics: a 12-year survey in elite athletes.

- Curr Opin Allergy Clin Immunol* (2015) 15(2):184–92. doi:10.1097/ACI.000000000000149
43. Bramley TJ, Lerner D, Sames M. Productivity losses related to the common cold. *J Occup Environ Med* (2002) 44(9):822–9. doi:10.1097/00043764-200209000-00004
 44. Fendrick AM, Monto AS, Nightengale B, Sarnes M. The economic burden of non-influenza-related viral respiratory tract infection in the United States. *Arch Intern Med* (2003) 163(4):487–94. doi:10.1001/archinte.163.4.487
 45. Bayer C, Remschmidt C, an der Heiden M, Tolksdorf K, Herzhoff M, Kaersten S, et al. Internet-based syndromic monitoring of acute respiratory illness in the general population of Germany, weeks 35/2011 to 34/2012. *Euro Surveill* (2014) 19(4):1–11. doi:10.2807/1560-7917.ES2014.19.4.20684
 46. Gleeson M, Pyne DB, Austin JP, Lynn Francis J, Clancy RL, McDonald WA, et al. Epstein-Barr virus reactivation and upper-respiratory illness in elite swimmers. *Med Sci Sports Exerc* (2002) 34(3):411–7. doi:10.1097/00005768-200203000-00005
 47. Cox AJ, Gleeson M, Pyne DB, Saunders PU, Clancy RL, Fricker PA. Valtrex therapy for Epstein-Barr virus reactivation and upper respiratory symptoms in elite runners. *Med Sci Sports Exerc* (2004) 36(7):1104–10. doi:10.1249/01.MSS.0000131957.40985.2B
 48. Speir E. Cytomegalovirus gene regulation by reactive oxygen species. Agents in atherosclerosis. *Ann N Y Acad Sci* (2000) 899:363–74. doi:10.1111/j.1749-6632.2000.tb06200.x
 49. Docke WD, Prosch S, Fietze E, Kimel V, Zuckermann H, Klug C, et al. Cytomegalovirus reactivation and tumour necrosis factor. *Lancet* (1994) 343(8892):268–9. doi:10.1016/S0140-6736(94)91116-9
 50. Prösch S, Wendt CEC, Reinke P, Priemer C, Oppert M, Krüger DH, et al. A novel link between stress and human cytomegalovirus (HCMV) infection: sympathetic hyperactivity stimulates HCMV activation. *Virology* (2000) 272:357–65. doi:10.1006/viro.2000.0367
 51. He CS, Handzlik M, Muhamad A, Gleeson M. Influence of CMV/EBV serostatus on respiratory infection incidence during 4 months of winter training in a student cohort of endurance athletes. *Eur J Appl Physiol* (2013) 113(10):2613–9. doi:10.1007/s00421-013-2704-x
 52. Bishop NC, Gleeson M. Acute and chronic effects of exercise on markers of mucosal immunity. *Front Biosci (Landmark Ed)* (2009) 14:4444–56. doi:10.2741/3540
 53. Tomasi TB, Trudeau FB, Czerwinski D, Erredge S. Immune parameters in athletes before and after strenuous exercise. *J Clin Immunol* (1982) 2(3):173–8. doi:10.1007/BF00915219
 54. Mackinnon LT, Chick TW, van As A, Tomasi TB. Decreased secretory immunoglobulins following intense endurance exercise. *Sports Train Med Rehabil* (1989) 1:209–18. doi:10.1080/15438628909511878
 55. Nieman DC, Henson DA, Fagoaga OR, Utter AC, Vinci DM, Davis JM, et al. Change in salivary IgA following a competitive marathon race. *Int J Sports Med* (2002) 23(1):69–75. doi:10.1055/s-2002-19375
 56. Nehlsen-Cannarella SL, Nieman DC, Fagoaga OR, Kelln WJ, Henson DA, Shannon M, et al. Saliva immunoglobulins in elite women rowers. *Eur J Appl Physiol* (2000) 81(3):222–8. doi:10.1007/s004210050034
 57. Palmer FM, Nieman DC, Henson DA, McAnulty SR, McAnulty L, Swick NS, et al. Influence of vitamin C supplementation on oxidative and salivary IgA changes following an ultramarathon. *Eur J Appl Physiol* (2003) 89(1):100–7. doi:10.1007/s00421-002-0756-4
 58. Steerenberg PA, van Asperen IA, van Nieuw Amerongen A, Biewenga A, Mol D, Medema GJ. Salivary levels of immunoglobulin A in triathletes. *Eur J Oral Sci* (1997) 105(4):305–9. doi:10.1111/j.1600-0722.1997.tb00245.x
 59. Nieman DC, Dumke CI, Henson DA, McAnulty SR, McAnulty LS, Lind RH, et al. Immune and oxidative changes during and following the western states endurance run. *Int J Sports Med* (2003) 24(7):541–7. doi:10.1055/s-2003-42018
 60. Blannin AK, Robson PJ, Walsh NP, Clark AM, Glennon L, Gleeson M. The effect of exercising to exhaustion at different intensities on saliva immunoglobulin A, protein and electrolyte secretion. *Int J Sports Med* (1998) 19(8):547–52. doi:10.1055/s-2007-971958
 61. Killer SC, Svendsen IS, Gleeson M. The influence of hydration status during prolonged endurance exercise on salivary antimicrobial proteins. *Eur J Appl Physiol* (2015) 115(9):1887–95. doi:10.1007/s00421-015-3173-1
 62. Allgrove JE, Gomes E, Hough J, Gleeson M. Effects of exercise intensity on salivary antimicrobial proteins and markers of stress in active men. *J Sports Sci* (2008) 26(6):653–61. doi:10.1080/02640410701716790
 63. Sari-Sarraf V, Reilly T, Doran DA. Salivary IgA response to intermittent and continuous exercise. *Int J Sports Med* (2006) 27(11):849–55. doi:10.1055/s-2006-923777
 64. Reid MR, Drummond PD, Mackinnon LT. The effect of moderate aerobic exercise and relaxation on secretory immunoglobulin A. *Int J Sports Med* (2001) 22(2):132–7. doi:10.1055/s-2001-11347
 65. Davison G. Innate immune responses to a single session of sprint interval training. *Appl Physiol Nutr Metab* (2011) 36(3):395–404. doi:10.1139/h11-033
 66. Walsh NP, Blannin AK, Clark AM, Cook L, Robson PJ, Gleeson M. The effects of high-intensity intermittent exercise on saliva IgA, total protein and alpha-amylase. *J Sports Sci* (1999) 17(2):129–34. doi:10.1080/026404199366226
 67. Peters EM, Shaik J, Kleinvelde N. Upper respiratory tract infection symptoms in ultramarathon runners not related to immunoglobulin status. *Clin J Sport Med* (2010) 20(1):39–46. doi:10.1097/JSM.0b013e3181cb4086
 68. Gleeson M, Bishop N, Oliveira M, McCauley T, Tauler P, Muhamad AS. Respiratory infection risk in athletes: association with antigen-stimulated IL-10 production and salivary IgA secretion. *Scand J Med Sci Sports* (2012) 22(3):410–7. doi:10.1111/j.1600-0838.2010.01272.x
 69. Gleeson M, Bishop N, Oliveira M, Tauler P. Influence of training load on upper respiratory tract infection incidence and antigen-stimulated cytokine production. *Scand J Med Sci Sports* (2013) 23(4):451–7. doi:10.1111/j.1600-0838.2011.01422.x
 70. Brandtzaeg P. Secretory immunity with special reference to the oral cavity. *J Oral Microbiol* (2013) 5:1–24. doi:10.3402/jom.v5i0.20401
 71. Marcotte H, Lavoie MC. Oral microbial ecology and the role of salivary immunoglobulin A. *Microbiol Mol Biol Rev* (1998) 62(1):71–109.
 72. Needleman I, Ashley P, Petrie A, Fortune F, Turner W, Jones J, et al. Oral health and impact on performance of athletes participating in the London 2012 Olympic Games: a cross-sectional study. *Br J Sports Med* (2013) 47(16):1054–8. doi:10.1136/bjsports-2013-092891
 73. Heaney JL, Gleeson M, Phillips AC, Taylor IM, Drayson MT, Goodall M, et al. Salivary immunoglobulin free light chains: reference ranges and responses to exercise in young and older adults. *Exerc Immunol Rev* (2016) 22:28–41.
 74. Li TL, Gleeson M. The effect of single and repeated bouts of prolonged cycling and circadian variation on saliva flow rate, immunoglobulin A and alpha-amylase responses. *J Sports Sci* (2004) 22(11–12):1015–24. doi:10.1080/02640410410001716733
 75. England CG, Hugo FN, Hilgert JB, Nascimento GG, Junges R, Lim HJ, et al. Psychological distress and salivary secretory immunity. *Brain Behav Immun* (2016) 52:11–7. doi:10.1016/j.bbi.2015.08.017
 76. Deslauriers N, Oudghiri M, Seguin J, Trudel L. The oral immune system: dynamics of salivary immunoglobulin production in the inbred mouse. *Immunol Invest* (1986) 15(4):339–49. doi:10.3109/08820138609052953
 77. Booth CK, Dwyer DB, Pacque PF, Ball MJ. Measurement of immunoglobulin A in saliva by particle-enhanced nephelometric immunoassay: sample collection, limits of quantitation, precision, stability and reference range. *Ann Clin Biochem* (2009) 46(Pt 5):401–6. doi:10.1258/acb.2009.008248
 78. Brandtzaeg P. Do salivary antibodies reliably reflect both mucosal and systemic immunity? *Ann N Y Acad Sci* (2007) 1098:288–311. doi:10.1196/annals.1384.012
 79. Hanstock HG, Walsh NP, Edwards JP, Fortes MB, Cosby SL, Nugent A, et al. Tear fluid SIgA as a noninvasive biomarker of mucosal immunity and common cold risk. *Med Sci Sports Exerc* (2016) 48(3):569–77. doi:10.1249/MSS.0000000000000801
 80. Gleeson M, Pyne DB, Elkington LJ, Hall ST, Attia JR, Oldmeadow C, et al. Developing a multi-component immune model for evaluating the risk of respiratory illness in athletes. *Exerc Immunol Rev* (2017) 23:52–64.
 81. Campbell JP, Riddell NE, Burns VE, Turner M, van Zanten JJ, Drayson MT, et al. Acute exercise mobilises CD8+ T lymphocytes exhibiting an effector-memory phenotype. *Brain Behav Immun* (2009) 23(6):767–75. doi:10.1016/j.bbi.2009.02.011
 82. Shephard RJ. Adhesion molecules, catecholamines and leucocyte redistribution during and following exercise. *Sports Med* (2003) 33(4):261–84. doi:10.2165/00007256-200333040-00002

83. Benschop RJ, Nijkamp FP, Ballieux RE, Heijnen CJ. The effects of beta-adrenoceptor stimulation on adhesion of human natural killer cells to cultured endothelium. *Br J Pharmacol* (1994) 113(4):1311–6. doi:10.1111/j.1476-5381.1994.tb17141.x
84. Kruger K, Lechtermann A, Fobker M, Volker K, Mooren FC. Exercise-induced redistribution of T lymphocytes is regulated by adrenergic mechanisms. *Brain Behav Immun* (2008) 22(3):324–38. doi:10.1016/j.bbi.2007.08.008
85. Dimitrov S, Lange T, Born J. Selective mobilization of cytotoxic leukocytes by epinephrine. *J Immunol* (2010) 184(1):503–11. doi:10.4049/jimmunol.0902189
86. Turner JE, Spielmann G, Wadley AJ, Aldred S, Simpson RJ, Campbell JP. Exercise-induced B cell mobilisation: preliminary evidence for an influx of immature cells into the bloodstream. *Physiol Behav* (2016) 164(Pt A):376–82. doi:10.1016/j.physbeh.2016.06.023
87. Kruger K, Alack K, Ringsels R, Mink L, Pfeifer E, Schinle M, et al. Apoptosis of T-cell subsets after acute high-intensity interval exercise. *Med Sci Sports Exerc* (2016) 48(10):2021–9. doi:10.1249/MSS.0000000000000979
88. Clifford T, Wood MJ, Stocks P, Howatson G, Stevenson EJ, Hilkens CMU. T-regulatory cells exhibit a biphasic response to prolonged endurance exercise in humans. *Eur J Appl Physiol* (2017) 117(8):1727–37. doi:10.1007/s00421-017-3667-0
89. Shek PN, Sabiston BH, Buguet A, Radomski MW. Strenuous exercise and immunological changes: a multiple-time-point analysis of leukocyte subsets, CD4/CD8 ratio, immunoglobulin production and NK cell response. *Int J Sports Med* (1995) 16(7):466–74. doi:10.1055/s-2007-973039
90. Shinkai S, Shore S, Shek PN, Shephard RJ. Acute exercise and immune function. Relationship between lymphocyte activity and changes in subset counts. *Int J Sports Med* (1992) 13(6):452–61. doi:10.1055/s-2007-1021297
91. Kakani MW, Peake J, Brenu EW, Simmonds M, Gray B, Hooper SL, et al. The open window of susceptibility to infection after acute exercise in healthy young male elite athletes. *Exerc Immunol Rev* (2010) 16:119–37.
92. Pedersen BK, Ullum H. NK cell response to physical activity: possible mechanisms of action. *Med Sci Sports Exerc* (1994) 26(2):140–6. doi:10.1249/00005768-199402000-00003
93. Dhabhar FS. Effects of stress on immune function: the good, the bad, and the beautiful. *Immunol Res* (2014) 58(2–3):193–210. doi:10.1007/s12026-014-8517-0
94. Kruger K, Mooren FC. T cell homing and exercise. *Exerc Immunol Rev* (2007) 13:37–54.
95. Mooren FC, Kruger K. Apoptotic lymphocytes induce progenitor cell mobilization after exercise. *J Appl Physiol* (1985) (2015) 119(2):135–9. doi:10.1152/jappphysiol.00287.2015
96. Mars M, Govender S, Weston A, Naicker V, Chuturgoon A. High intensity exercise: a cause of lymphocyte apoptosis? *Biochem Biophys Res Commun* (1998) 249(2):366–70. doi:10.1006/bbrc.1998.9156
97. Mooren FC, Lechtermann A, Volker K. Exercise-induced apoptosis of lymphocytes depends on training status. *Med Sci Sports Exerc* (2004) 36(9):1476–83. doi:10.1249/01.MSS.0000139897.34521.E9
98. Mooren FC, Blomling D, Lechtermann A, Lerch MM, Volker K. Lymphocyte apoptosis after exhaustive and moderate exercise. *J Appl Physiol* (2002) 93(1):147–53. doi:10.1152/jappphysiol.01262.2001
99. Tanimura Y, Shimizu K, Tanabe K, Otsuki T, Yamauchi R, Matsubara Y, et al. Exercise-induced oxidative DNA damage and lymphocytopenia in sedentary young males. *Med Sci Sports Exerc* (2008) 40(8):1455–62. doi:10.1249/MSS.0b013e31817242cf
100. Simpson RJ, Florida-James GD, Whyte GP, Black JR, Ross JA, Guy K. Apoptosis does not contribute to the blood lymphocytopenia observed after intensive and downhill treadmill running in humans. *Res Sports Med* (2007) 15(3):157–74. doi:10.1080/15438620701405339
101. Shephard RJ, Shek PN. Effects of exercise and training on natural killer cell counts and cytolytic activity: a meta-analysis. *Sports Med* (1999) 28(3):177–95. doi:10.2165/00007256-199928030-00003
102. Hansen JB, Wilsbard L, Osterud B. Biphasic changes in leukocytes induced by strenuous exercise. *Eur J Appl Physiol Occup Physiol* (1991) 62(3):157–61. doi:10.1007/BF00643735
103. Freud AG, Caligiuri MA. Human natural killer cell development. *Immunol Rev* (2006) 214:56–72. doi:10.1111/j.1600-065X.2006.00451.x
104. Poli A, Michel T, Theresine M, Andres E, Hentges F, Zimmer J. CD56bright natural killer (NK) cells: an important NK cell subset. *Immunology* (2009) 126(4):458–65. doi:10.1111/j.1365-2567.2008.03027.x
105. Bjorkstrom NK, Riese P, Heuts F, Andersson S, Fauriat C, Ivarsson MA, et al. Expression patterns of NKG2A, KIR, and CD57 define a process of CD56dim NK-cell differentiation uncoupled from NK-cell education. *Blood* (2010) 116(19):3853–64. doi:10.1182/blood-2010-04-281675
106. Lopez-Verges S, Milush JM, Pandey S, York VA, Arakawa-Hoyt J, Pircher H, et al. CD57 defines a functionally distinct population of mature NK cells in the human CD56dimCD16+ NK-cell subset. *Blood* (2010) 116(19):3865–74. doi:10.1182/blood-2010-04-282301
107. Bigley AB, Rezvani K, Chew C, Sekine T, Pistillo M, Crucian B, et al. Acute exercise preferentially redeploys NK-cells with a highly-differentiated phenotype and augments cytotoxicity against lymphoma and multiple myeloma target cells. *Brain Behav Immun* (2014) 39:160–71. doi:10.1016/j.bbi.2013.10.030
108. Bigley AB, Rezvani K, Pistillo M, Reed J, Agha N, Kunz H, et al. Acute exercise preferentially redeploys NK-cells with a highly-differentiated phenotype and augments cytotoxicity against lymphoma and multiple myeloma target cells. Part II: impact of latent cytomegalovirus infection and catecholamine sensitivity. *Brain Behav Immun* (2015) 49:59–65. doi:10.1016/j.bbi.2014.12.027
109. Simpson RJ, Florida-James GD, Cosgrove C, Whyte GP, Macrae S, Pircher H, et al. High-intensity exercise elicits the mobilization of senescent T lymphocytes into the peripheral blood compartment in human subjects. *J Appl Physiol* (1985) (2007) 103(1):396–401. doi:10.1152/jappphysiol.00007.2007
110. Simpson RJ, Cosgrove C, Ingram LA, Florida-James GD, Whyte GP, Pircher H, et al. Senescent T-lymphocytes are mobilised into the peripheral blood compartment in young and older humans after exhaustive exercise. *Brain Behav Immun* (2008) 22(4):544–51. doi:10.1016/j.bbi.2007.11.002
111. Simpson RJ, Cosgrove C, Chee MM, McFarlin BK, Bartlett DB, Spielmann G, et al. Senescent phenotypes and telomere lengths of peripheral blood T-cells mobilized by acute exercise in humans. *Exerc Immunol Rev* (2010) 16:40–55.
112. Pedersen L, Idorn M, Olofsson GH, Lauenborg B, Nookaew I, Hansen RH, et al. Voluntary running suppresses tumor growth through epinephrine- and IL-6-dependent NK cell mobilization and redistribution. *Cell Metab* (2016) 23(3):554–62. doi:10.1016/j.cmet.2016.01.011
113. Kim R, Emi M, Tanabe K. Cancer immunoediting from immune surveillance to immune escape. *Immunology* (2007) 121(1):1–14. doi:10.1111/j.1365-2567.2007.02587.x
114. Scheiermann C, Kunisaki Y, Frenette PS. Circadian control of the immune system. *Nat Rev Immunol* (2013) 13(3):190–8. doi:10.1038/nri3386
115. Suzuki K, Hayano Y, Nakai A, Furuta F, Noda M. Adrenergic control of the adaptive immune response by diurnal lymphocyte recirculation through lymph nodes. *J Exp Med* (2016) 213(12):2567–74. doi:10.1084/jem.20160723
116. Dimitrov S, Benedict C, Heutling D, Westermann J, Born J, Lange T. Cortisol and epinephrine control opposing circadian rhythms in T cell subsets. *Blood* (2009) 113(21):5134–43. doi:10.1182/blood-2008-11-190769
117. Hojman P. Exercise protects from cancer through regulation of immune function and inflammation. *Biochem Soc Trans* (2017) 45(4):905–11. doi:10.1042/BST20160466
118. Turner JE, Wadley AJ, Aldred S, Fisher JP, Bosch JA, Campbell JP. Intensive exercise does not preferentially mobilize skin-homing T cells and NK cells. *Med Sci Sports Exerc* (2016) 48(7):1285–93. doi:10.1249/MSS.0000000000000914
119. Hanson ED, Danson E, Nguyen-Robertson CV, Fyfe JJ, Stepto NK, Bartlett DB, et al. Maximal exercise increases mucosal associated invariant T cell frequency and number in healthy young men. *Eur J Appl Physiol* (2017) 117(11):2159–69. doi:10.1007/s00421-017-3704-z
120. Gross E, Sunwoo JB, Bui JD. Cancer immunosurveillance and immunoediting by natural killer cells. *Cancer J* (2013) 19(6):483–9. doi:10.1097/PPO.0000000000000005
121. Simpson RJ, Bigley AB, Agha N, Hanley PJ, Bollard CM. Mobilizing immune cells with exercise for cancer immunotherapy. *Exerc Sport Sci Rev* (2017) 45(3):163–72. doi:10.1249/JES.0000000000000114
122. Moore SC, Lee IM, Weiderpass E, Campbell PT, Sampson JN, Kitahara CM, et al. Association of leisure-time physical activity with risk of 26 types of

- cancer in 1.44 million adults. *JAMA Intern Med* (2016) 176(6):816–25. doi:10.1001/jamainternmed.2016.1548
123. Turner JE, Aldred S, Witard OC, Drayson MT, Moss PM, Bosch JA. Latent cytomegalovirus infection amplifies CD8 T-lymphocyte mobilisation and egress in response to exercise. *Brain Behav Immun* (2010) 24(8):1362–70. doi:10.1016/j.bbi.2010.07.239
 124. Lancaster GI, Khan Q, Drysdale PT, Wallace F, Jeukendrup AE, Drayson MT, et al. Effect of prolonged exercise and carbohydrate ingestion on type 1 and type 2 T lymphocyte distribution and intracellular cytokine production in humans. *J Appl Physiol* (1985) (2005) 98(2):565–71. doi:10.1152/jappphysiol.00754.2004
 125. Starkie RL, Rolland J, Febbraio MA. Effect of adrenergic blockade on lymphocyte cytokine production at rest and during exercise. *Am J Physiol Cell Physiol* (2001) 281(4):C1233–40. doi:10.1152/ajpcell.2001.281.4.C1233
 126. Siedlik JA, Deckert JA, Benedict SH, Bhatta A, Dunbar AJ, Vardiman JP, et al. T cell activation and proliferation following acute exercise in human subjects is altered by storage conditions and mitogen selection. *J Immunol Methods* (2017) 446:7–14. doi:10.1016/j.jim.2017.03.017
 127. Siedlik JA, Benedict SH, Landes EJ, Weir JP, Vardiman JP, Gallagher PM. Acute bouts of exercise induce a suppressive effect on lymphocyte proliferation in human subjects: a meta-analysis. *Brain Behav Immun* (2016) 56:343–51. doi:10.1016/j.bbi.2016.04.008
 128. Zimmer P, Schenk A, Kieven M, Holthaus M, Lehmann J, Lovenich L, et al. Exercise induced alterations in NK-cell cytotoxicity – methodological issues and future perspectives. *Exerc Immunol Rev* (2017) 23:66–81.
 129. McFarlin BK, Venable AS, Carpenter CK, Henning AL, Stephan O. Oral supplementation with baker's yeast beta glucan is associated with altered monocytes, T cells and cytokines following a bout of strenuous exercise. *Front Physiol* (2017) 8:786. doi:10.3389/fphys.2017.00786
 130. Witard OC, Turner JE, Jackman SR, Kies AK, Jeukendrup AE, Bosch JA, et al. High dietary protein restores overreaching induced impairments in leukocyte trafficking and reduces the incidence of upper respiratory tract infection in elite cyclists. *Brain Behav Immun* (2014) 39:211–9. doi:10.1016/j.bbi.2013.10.002
 131. Albers R, Bourdet-Sicard R, Braun D, Calder PC, Herz U, Lambert C, et al. Monitoring immune modulation by nutrition in the general population: identifying and substantiating effects on human health. *Br J Nutr* (2013) 110(Suppl 2):S1–30. doi:10.1017/S0007114513001505
 132. Burns VE, Gallagher S. Antibody response to vaccination as a marker of in vivo immune function in psychophysiological research. *Neurosci Biobehav Rev* (2010) 35(1):122–6. doi:10.1016/j.neubiorev.2010.01.005
 133. Edwards KM, Burns VE, Reynolds T, Carroll D, Drayson M, Ring C. Acute stress exposure prior to influenza vaccination enhances antibody response in women. *Brain Behav Immun* (2006) 20(2):159–68. doi:10.1016/j.bbi.2005.07.001
 134. Edwards KM, Burns VE, Allen LM, McPhee JS, Bosch JA, Carroll D, et al. Eccentric exercise as an adjuvant to influenza vaccination in humans. *Brain Behav Immun* (2007) 21(2):209–17. doi:10.1016/j.bbi.2006.04.158
 135. Edwards KM, Burns VE, Adkins AE, Carroll D, Drayson M, Ring C. Meningococcal A vaccination response is enhanced by acute stress in men. *Psychosom Med* (2008) 70(2):147–51. doi:10.1097/PSY.0b013e318164232e
 136. Edwards KM, Campbell JP, Ring C, Drayson MT, Bosch JA, Downes C, et al. Exercise intensity does not influence the efficacy of eccentric exercise as a behavioural adjuvant to vaccination. *Brain Behav Immun* (2010) 24(4):623–30. doi:10.1016/j.bbi.2010.01.009
 137. Edwards KM, Pung MA, Tomfohr LM, Ziegler MG, Campbell JP, Drayson MT, et al. Acute exercise enhancement of pneumococcal vaccination response: a randomised controlled trial of weaker and stronger immune response. *Vaccine* (2012) 30(45):6389–95. doi:10.1016/j.vaccine.2012.08.022
 138. Ranadive SM, Cook M, Kappus RM, Yan H, Lane AD, Woods JA, et al. Effect of acute aerobic exercise on vaccine efficacy in older adults. *Med Sci Sports Exerc* (2014) 46(3):455–61. doi:10.1249/MSS.0b013e3182a75ff2
 139. Long JE, Ring C, Drayson M, Bosch J, Campbell JP, Bhabra J, et al. Vaccination response following aerobic exercise: can a brisk walk enhance antibody response to pneumococcal and influenza vaccinations? *Brain Behav Immun* (2012) 26(4):680–7. doi:10.1016/j.bbi.2012.02.004
 140. Edwards KM, Pascoe AR, Fiatarone-Singh MA, Singh NA, Kok J, Booy R. A randomised controlled trial of resistance exercise prior to administration of influenza vaccination in older adults. *Brain Behav Immun* (2015) 49:e24–5. doi:10.1016/j.bbi.2015.06.102
 141. Furman D, Hejblum BP, Simon N, Jojic V, Dekker CL, Thiebaut R, et al. Systems analysis of sex differences reveals an immunosuppressive role for testosterone in the response to influenza vaccination. *Proc Natl Acad Sci U S A* (2014) 111(2):869–74. doi:10.1073/pnas.1321060111
 142. Trigunaita A, Dimo J, Jorgensen TN. Suppressive effects of androgens on the immune system. *Cell Immunol* (2015) 294(2):87–94. doi:10.1016/j.cellimm.2015.02.004
 143. Haq K, McElhaney JE. Immunosenescence: influenza vaccination and the elderly. *Curr Opin Immunol* (2014) 29:38–42. doi:10.1016/j.coi.2014.03.008
 144. Kohut ML, Cooper MM, Nickolaus MS, Russell DR, Cunnick JE. Exercise and psychosocial factors modulate immunity to influenza vaccine in elderly individuals. *J Gerontol A Biol Sci Med Sci* (2002) 57(9):M557–62. doi:10.1093/gerona/57.9.M557
 145. Kohut ML, Arntson BA, Lee W, Rozeboom K, Yoon KJ, Cunnick JE, et al. Moderate exercise improves antibody response to influenza immunization in older adults. *Vaccine* (2004) 22(17–18):2298–306. doi:10.1016/j.vaccine.2003.11.023
 146. Smith TP, Kennedy SL, Fleshner M. Influence of age and physical activity on the primary in vivo antibody and T cell-mediated responses in men. *J Appl Physiol* (1985) (2004) 97(2):491–8. doi:10.1152/jappphysiol.01404.2003
 147. Schuler PB, Leblanc PA, Marzilli TS. Effect of physical activity on the production of specific antibody in response to the 1998–99 influenza virus vaccine in older adults. *J Sports Med Phys Fitness* (2003) 43(3):404.
 148. Grant RW, Mariani RA, Vieira VJ, Fleshner M, Smith TP, Keylock KT, et al. Cardiovascular exercise intervention improves the primary antibody response to keyhole limpet hemocyanin (KLH) in previously sedentary older adults. *Brain Behav Immun* (2008) 22(6):923–32. doi:10.1016/j.bbi.2008.01.006
 149. Woods JA, Keylock KT, Lowder T, Vieira VJ, Zerkovich W, Dumich S, et al. Cardiovascular exercise training extends influenza vaccine seroprotection in sedentary older adults: the immune function intervention trial. *J Am Geriatr Soc* (2009) 57(12):2183–91. doi:10.1111/j.1532-5415.2009.02563.x
 150. Yang Y, Verkuilen J, Rosengren KS, Mariani RA, Reed M, Grubisich SA, et al. Effects of a traditional Taiji/Qigong curriculum on older adults' immune response to influenza vaccine. *Med Sport Sci* (2008) 52:64–76. doi:10.1159/000134285
 151. de Araujo AL, Silva LC, Fernandes JR, Matias Mde S, Boas LS, Machado CM, et al. Elderly men with moderate and intense training lifestyle present sustained higher antibody responses to influenza vaccine. *Age (Dordr)* (2015) 37(6):105. doi:10.1007/s11357-015-9843-4
 152. Kapasi ZF, Catlin PA, Joyner DR, Lewis ML, Schwartz AL, Townsend EL. The effects of intense physical exercise on secondary antibody response in young and old mice. *Phys Ther* (2000) 80(11):1076–86.
 153. Grande AJ, Reid H, Thomas EE, Nunan D, Foster C. Exercise prior to influenza vaccination for limiting influenza incidence and its related complications in adults. *Cochrane Database Syst Rev* (2016) (8):CD011857. doi:10.1002/14651858.CD011857.pub2
 154. Warren K, Thompson N, Wannemuehler M, Kohut M. Antibody and CD8+ T cell memory response to influenza A/PR/8/34 infection is reduced in treadmill-exercised mice, yet still protective. *J Appl Physiol* (1985) (2013) 114(10):1413–20. doi:10.1152/jappphysiol.01355.2012
 155. Sim YJ, Yu S, Yoon KJ, Loiacono CM, Kohut ML. Chronic exercise reduces illness severity, decreases viral load, and results in greater anti-inflammatory effects than acute exercise during influenza infection. *J Infect Dis* (2009) 200(9):1434–42. doi:10.1086/606014
 156. Lowder T, Padgett DA, Woods JA. Moderate exercise protects mice from death due to influenza virus. *Brain Behav Immun* (2005) 19(5):377–80. doi:10.1016/j.bbi.2005.04.002
 157. Nosbaum A, Vocanson M, Rozieres A, Hennino A, Nicolas JF. Allergic and irritant contact dermatitis. *Eur J Dermatol* (2009) 19(4):325–32. doi:10.1684/ejd.2009.0686
 158. Gulati N, Suarez-Farinas M, Fuentes-Duculan J, Gilleaudeau P, Sullivan-Whalen M, Correa da Rosa J, et al. Molecular characterization of human skin response to diphencyprone at peak and resolution phases: therapeutic insights. *J Invest Dermatol* (2014) 134(10):2531–40. doi:10.1038/jid.2014.196

159. Diment BC, Fortes MB, Edwards JP, Hanstock HG, Ward MD, Dunstall HM, et al. Exercise intensity and duration effects on in vivo immunity. *Med Sci Sports Exerc* (2015) 47(7):1390–8. doi:10.1249/MSS.0000000000000562
160. Harper Smith AD, Coakley SL, Ward MD, Macfarlane AW, Friedmann PS, Walsh NP. Exercise-induced stress inhibits both the induction and elicitation phases of in vivo T-cell-mediated immune responses in humans. *Brain Behav Immun* (2011) 25(6):1136–42. doi:10.1016/j.bbi.2011.02.014
161. Ceddia MA, Price EA, Kohlmeier CK, Evans JK, Lu Q, McAuley E, et al. Differential leukocytosis and lymphocyte mitogenic response to acute maximal exercise in the young and old. *Med Sci Sports Exerc* (1999) 31(6):829–36. doi:10.1097/00005768-199906000-00011
162. Woods JA, Evans JK, Wolters BW, Ceddia MA, McAuley E. Effects of maximal exercise on natural killer (NK) cell cytotoxicity and responsiveness to interferon-alpha in the young and old. *J Gerontol A Biol Sci Med Sci* (1998) 53(6):B430–7. doi:10.1093/gerona/53A.6.B430
163. Simpson RJ, Lowder TW, Spielmann G, Bigley AB, LaVoy EC, Kunz H. Exercise and the aging immune system. *Ageing Res Rev* (2012) 11(3):404–20. doi:10.1016/j.arr.2012.03.003
164. Kohut ML, Senchina DS. Reversing age-associated immunosenescence via exercise. *Exerc Immunol Rev* (2004) 10:6–41.
165. Woods JA, Lowder TW, Keylock KT. Can exercise training improve immune function in the aged? *Ann N Y Acad Sci* (2002) 959:117–27. doi:10.1111/j.1749-6632.2002.tb02088.x
166. Simpson RJ, Bigley AB, Spielmann G, LaVoy EC, Kunz H, Bollard CM. Human cytomegalovirus infection and the immune response to exercise. *Exerc Immunol Rev* (2016) 22:8–27.
167. Bartlett DB, Fox O, McNulty CL, Greenwood HL, Murphy L, Sapey E, et al. Habitual physical activity is associated with the maintenance of neutrophil migratory dynamics in healthy older adults. *Brain Behav Immun* (2016) 56:12–20. doi:10.1016/j.bbi.2016.02.024
168. Bartlett DB, Shepherd SO, Wilson OJ, Adlan AM, Wagenmakers AJM, Shaw CS, et al. Neutrophil and monocyte bactericidal responses to 10 weeks of low-volume high-intensity interval or moderate-intensity continuous training in sedentary adults. *Oxid Med Cell Longev* (2017) 2017:8148742. doi:10.1155/2017/8148742
169. Shephard RJ. Development of the discipline of exercise immunology. *Exerc Immunol Rev* (2010) 16:194–222.
170. Hazeldine J, Lord JM. Innate immunosenescence: underlying mechanisms and clinical relevance. *Biogerontology* (2015) 16(2):187–201. doi:10.1007/s10522-014-9514-3
171. Peralbo E, Alonso C, Solana R. Invariant NKT and NKT-like lymphocytes: two different T cell subsets that are differentially affected by ageing. *Exp Gerontol* (2007) 42(8):703–8. doi:10.1016/j.exger.2007.05.002
172. Ziegler-Heitbrock L, Ancuta P, Crowe S, Dalod M, Grau V, Hart DN, et al. Nomenclature of monocytes and dendritic cells in blood. *Blood* (2010) 116(16):e74–80. doi:10.1182/blood-2010-02-258558
173. Goh J, Ladiges WC. Exercise enhances wound healing and prevents cancer progression during aging by targeting macrophage polarity. *Mech Ageing Dev* (2014) 139:41–8. doi:10.1016/j.mad.2014.06.004
174. Kanneganti TD, Dixit VD. Immunological complications of obesity. *Nat Immunol* (2012) 13(8):707–12. doi:10.1038/ni.2343
175. Garg SK, Delaney C, Shi H, Yung R. Changes in adipose tissue macrophages and T cells during aging. *Crit Rev Immunol* (2014) 34(1):1–14. doi:10.1615/CritRevImmunol.2013006833
176. Kratz M, Coats BR, Hisert KB, Hagman D, Mutskov V, Peris E, et al. Metabolic dysfunction drives a mechanistically distinct proinflammatory phenotype in adipose tissue macrophages. *Cell Metab* (2014) 20(4):614–25. doi:10.1016/j.cmet.2014.08.010
177. Bourlier V, Zakaroff-Girard A, Miranville A, De Barros S, Maumus M, Sengenès C, et al. Remodeling phenotype of human subcutaneous adipose tissue macrophages. *Circulation* (2008) 117(6):806–15. doi:10.1161/CIRCULATIONAHA.107.724096
178. Zeyda M, Farmer D, Todoric J, Aszmann O, Speiser M, Gyori G, et al. Human adipose tissue macrophages are of an anti-inflammatory phenotype but capable of excessive pro-inflammatory mediator production. *Int J Obes (Lond)* (2007) 31(9):1420–8. doi:10.1038/sj.ijo.0803632
179. Xu X, Grijalva A, Skowronski A, van Eijk M, Serlie MJ, Ferrante AW Jr. Obesity activates a program of lysosomal-dependent lipid metabolism in adipose tissue macrophages independently of classic activation. *Cell Metab* (2013) 18(6):816–30. doi:10.1016/j.cmet.2013.11.001
180. Chavez-Galan L, Olleros ML, Vesin D, Garcia I. Much more than M1 and M2 macrophages, there are also CD169(+) and TCR(+) macrophages. *Front Immunol* (2015) 6:263. doi:10.3389/fimmu.2015.00263
181. Mills CD. Anatomy of a discovery: m1 and m2 macrophages. *Front Immunol* (2015) 6:212. doi:10.3389/fimmu.2015.00212
182. Bueno V, Sant'Anna OA, Lord JM. Ageing and myeloid-derived suppressor cells: possible involvement in immunosenescence and age-related disease. *Age (Dordr)* (2014) 36(6):9729. doi:10.1007/s11357-014-9729-x
183. Shearer GM. Th1/Th2 changes in aging. *Mech Ageing Dev* (1997) 94(1–3):1–5.
184. Schmitt V, Rink L, Uciechowski P. The Th17/Treg balance is disturbed during aging. *Exp Gerontol* (2013) 48(12):1379–86. doi:10.1016/j.exger.2013.09.003
185. Fulop T, Larbi A, Pawelec G. Human T cell aging and the impact of persistent viral infections. *Front Immunol* (2013) 4:271. doi:10.3389/fimmu.2013.00271
186. Pawelec G. Hallmarks of human “immunosenescence”: adaptation or dysregulation? *Immun Ageing* (2012) 9(1):15. doi:10.1186/1742-4933-9-15
187. Geiger H, de Haan G, Florian MC. The ageing haematopoietic stem cell compartment. *Nat Rev Immunol* (2013) 13(5):376–89. doi:10.1038/nri3433
188. Roux A, Mourin G, Larsen M, Fastenackels S, Urrutia A, Gorochov G, et al. Differential impact of age and cytomegalovirus infection on the gamma-delta T cell compartment. *J Immunol* (2013) 191(3):1300–6. doi:10.4049/jimmunol.1202940
189. Colonna-Romano G, Aquino A, Bulati M, Lio D, Candore G, Oddo G, et al. Impairment of gamma/delta T lymphocytes in elderly: implications for immunosenescence. *Exp Gerontol* (2004) 39(10):1439–46. doi:10.1016/j.exger.2004.07.005
190. Michishita Y, Hirokawa M, Guo YM, Abe Y, Liu J, Ubukawa K, et al. Age-associated alteration of gammadelta T-cell repertoire and different profiles of activation-induced death of Vdelta1 and Vdelta2 T cells. *Int J Hematol* (2011) 94(3):230–40. doi:10.1007/s12185-011-0907-7
191. Jagger A, Shimojima Y, Goronzy JJ, Weyand CM. Regulatory T cells and the immune aging process: a mini-review. *Gerontology* (2014) 60(2):130–7. doi:10.1159/000355303
192. Siegrist CA, Aspinall R. B-cell responses to vaccination at the extremes of age. *Nat Rev Immunol* (2009) 9(3):185–94. doi:10.1038/nri2508
193. Appay V, Sauce D. Naive T cells: the crux of cellular immune aging? *Exp Gerontol* (2014) 54:90–3. doi:10.1016/j.exger.2014.01.003
194. Pawelec G. Immunosenescence: role of cytomegalovirus. *Exp Gerontol* (2013) 54:1–5. doi:10.1016/j.exger.2013.11.010
195. Wikby A, Maxson P, Olsson J, Johansson B, Ferguson FG. Changes in CD8 and CD4 lymphocyte subsets, T cell proliferation responses and non-survival in the very old: the Swedish longitudinal OCTO-immune study. *Mech Ageing Dev* (1998) 102(2–3):187–98. doi:10.1016/S0047-6374(97)00151-6
196. Pawelec G, Ferguson FG, Wikby A. The SENIEUR protocol after 16 years. *Mech Ageing Dev* (2001) 122(2):132–4. doi:10.1016/S0047-6374(00)00240-2
197. Wikby A, Johansson B, Ferguson F, Olsson J. Age-related changes in immune parameters in a very old population of Swedish people: a longitudinal study. *Exp Gerontol* (1994) 29(5):531–41. doi:10.1016/0531-5565(94)90036-1
198. Wikby A, Nilsson BO, Forsey R, Thompson J, Strindhall J, Lofgren S, et al. The immune risk phenotype is associated with IL-6 in the terminal decline stage: findings from the Swedish NONA immune longitudinal study of very late life functioning. *Mech Ageing Dev* (2006) 127(8):695–704. doi:10.1016/j.mad.2006.04.003
199. Wikby A, Ferguson F, Forsey R, Thompson J, Strindhall J, Lofgren S, et al. An immune risk phenotype, cognitive impairment, and survival in very late life: impact of allostatic load in Swedish octogenarian and nonagenarian humans. *J Gerontol A Biol Sci Med Sci* (2005) 60(5):556–65. doi:10.1093/gerona/60.5.556
200. Olsson J, Wikby A, Johansson B, Lofgren S, Nilsson BO, Ferguson FG. Age-related change in peripheral blood T-lymphocyte subpopulations and cytomegalovirus infection in the very old: the Swedish longitudinal OCTO immune study. *Mech Ageing Dev* (2000) 121(1–3):187–201. doi:10.1016/S0047-6374(00)00210-4
201. Franceschi C, Bonafe M, Valensin S, Olivieri F, De Luca M, Ottaviani E, et al. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci* (2000) 908:244–54. doi:10.1111/j.1749-6632.2000.tb06651.x

202. Franceschi C, Capri M, Monti D, Giunta S, Olivieri F, Sevini F, et al. Inflammaging and anti-inflammaging: a systemic perspective on aging and longevity emerged from studies in humans. *Mech Ageing Dev* (2007) 128(1):92–105. doi:10.1016/j.mad.2006.11.016
203. Giovannini S, Onder G, Liperoti R, Russo A, Carter C, Capoluongo E, et al. Interleukin-6, C-reactive protein, and tumor necrosis factor-alpha as predictors of mortality in frail, community-living elderly individuals. *J Am Geriatr Soc* (2011) 59(9):1679–85. doi:10.1111/j.1532-5415.2011.03570.x
204. Baune BT, Rothermundt M, Ladwig KH, Meisinger C, Berger K. Systemic inflammation (interleukin 6) predicts all-cause mortality in men: results from a 9-year follow-up of the MEMO Study. *Age (Dordr)* (2011) 33(2):209–17. doi:10.1007/s11357-010-9165-5
205. Bruunsgaard H, Ladelund S, Pedersen AN, Schroll M, Jorgensen T, Pedersen BK. Predicting death from tumour necrosis factor-alpha and interleukin-6 in 80-year-old people. *Clin Exp Immunol* (2003) 132(1):24–31. doi:10.1046/j.1365-2249.2003.02137.x
206. McElhaney JE, Kuchel GA, Zhou X, Swain SL, Haynes L. T-cell immunity to influenza in older adults: a pathophysiological framework for development of more effective vaccines. *Front Immunol* (2016) 7:41. doi:10.3389/fimmu.2016.00041
207. Frasca D, Blomberg BB. Aging, cytomegalovirus (CMV) and influenza vaccine responses. *Hum Vaccin Immunother* (2016) 12(3):682–90. doi:10.1080/21645515.2015.1105413
208. Brown FF, Bigley AB, Sherry C, Neal CM, Witard OC, Simpson RJ, et al. Training status and sex influence on senescent T-lymphocyte redistribution in response to acute maximal exercise. *Brain Behav Immun* (2014) 39:152–9. doi:10.1016/j.bbi.2013.10.031
209. Brown FF, Bigley AB, Ross JC, LaVoy EC, Simpson RJ, Galloway SD. T-lymphocyte populations following a period of high volume training in female soccer players. *Physiol Behav* (2015) 152(Pt A):175–81. doi:10.1016/j.physbeh.2015.09.027
210. Gustafson MP, DiCostanzo AC, Wheatley CM, Kim CH, Bornschlegl S, Gastineau DA, et al. A systems biology approach to investigating the influence of exercise and fitness on the composition of leukocytes in peripheral blood. *J Immunother Cancer* (2017) 5:30. doi:10.1186/s40425-017-0231-8
211. Prieto-Hinojosa A, Knight A, Compton C, Gleeson M, Travers PJ. Reduced thymic output in elite athletes. *Brain Behav Immun* (2014) 39:75–9. doi:10.1016/j.bbi.2014.01.004
212. Moro-Garcia MA, Fernandez-Garcia B, Echeverria A, Rodriguez-Alonso M, Suarez-Garcia FM, Solano-Jaurrieta JJ, et al. Frequent participation in high volume exercise throughout life is associated with a more differentiated adaptive immune response. *Brain Behav Immun* (2014) 39:61–74. doi:10.1016/j.bbi.2013.12.014
213. Teixeira AM, Rama L, Carvalho HM, Borges G, Carvalheiro T, Gleeson M, et al. Changes in naive and memory T-cells in elite swimmers during a winter training season. *Brain Behav Immun* (2014) 39:186–93. doi:10.1016/j.bbi.2014.01.002
214. Cosgrove C, Galloway SD, Neal C, Hunter AM, McFarlin BK, Spielmann G, et al. The impact of 6-month training preparation for an Ironman triathlon on the proportions of naive, memory and senescent T cells in resting blood. *Eur J Appl Physiol* (2012) 112(8):2989–98. doi:10.1007/s00421-011-2273-9
215. Dopic XC, Evangelou M, Ferreira RC, Guo H, Pekalski ML, Smyth DJ, et al. Widespread seasonal gene expression reveals annual differences in human immunity and physiology. *Nat Commun* (2015) 6:7000. doi:10.1038/ncomms8000
216. Goldinger A, Shakhbazov K, Henders AK, McRae AF, Montgomery GW, Powell JE. Seasonal effects on gene expression. *PLoS One* (2015) 10(5):e0126995. doi:10.1371/journal.pone.0126995
217. Spielmann G, McFarlin BK, O'Connor DP, Smith PJ, Pircher H, Simpson RJ. Aerobic fitness is associated with lower proportions of senescent blood T-cells in man. *Brain Behav Immun* (2011) 25(8):1521–9. doi:10.1016/j.bbi.2011.07.226
218. Silva LC, de Araujo AL, Fernandes JR, Matias Mde S, Silva PR, Duarte AJ, et al. Moderate and intense exercise lifestyles attenuate the effects of aging on telomere length and the survival and composition of T cell subpopulations. *Age (Dordr)* (2016) 38(1):24. doi:10.1007/s11357-016-9879-0
219. Duggal NA, Pollock RD, Lazarus NR, Harridge S, Lord JM. Major features of immunosenescence, including reduced thymic output, are ameliorated by high levels of physical activity in adulthood. *Aging Cell* (2018). doi:10.1111/acel.12750
220. Shimizu K, Kimura F, Akimoto T, Akama T, Tanabe K, Nishijima T, et al. Effect of moderate exercise training on T-helper cell subpopulations in elderly people. *Exerc Immunol Rev* (2008) 14:24–37.
221. Cao Dinh H, Beyer I, Mets T, Onyema OO, Njemini R, Renmans W, et al. Effects of physical exercise on markers of cellular immunosenescence: a systematic review. *Calcif Tissue Int* (2017) 100(2):193–215. doi:10.1007/s00223-016-0212-9
222. Kruger K, Frost S, Most E, Volker K, Pallauf J, Mooren FC. Exercise affects tissue lymphocyte apoptosis via redox-sensitive and Fas-dependent signaling pathways. *Am J Physiol Regul Integr Comp Physiol* (2009) 296(5):R1518–27. doi:10.1152/ajpregu.90994.2008
223. Khan N, Shariff N, Cobbold M, Bruton R, Ainsworth JA, Sinclair AJ, et al. Cytomegalovirus seropositivity drives the CD8 T cell repertoire toward greater clonality in healthy elderly individuals. *J Immunol* (2002) 169(4):1984–92. doi:10.4049/jimmunol.169.4.1984
224. Mueller YM, De Rosa SC, Hutton JA, Witek J, Roederer M, Altman JD, et al. Increased CD95/Fas-induced apoptosis of HIV-specific CD8(+) T cells. *Immunity* (2001) 15(6):871–82. doi:10.1016/S1074-7613(01)00246-1
225. Wang JS, Lin CT. Systemic hypoxia promotes lymphocyte apoptosis induced by oxidative stress during moderate exercise. *Eur J Appl Physiol* (2010) 108(2):371–82. doi:10.1007/s00421-009-1231-2
226. Takahashi A, Hanson MG, Norell HR, Havelka AM, Kono K, Malmberg KJ, et al. Preferential cell death of CD8+ effector memory (CCR7-CD45RA-) T cells by hydrogen peroxide-induced oxidative stress. *J Immunol* (2005) 174(10):6080–7. doi:10.4049/jimmunol.174.10.6080
227. Stocchi R, Ward KN, Fanin R, Bacarani M, Apperley JF. Management of human cytomegalovirus infection and disease after allogeneic bone marrow transplantation. *Haematologica* (1999) 84(1):71–9.
228. Derhovanessian E, Maier AB, Hahnel K, Zelba H, de Craen AJ, Roelofs H, et al. Lower proportion of naive peripheral CD8+ T cells and an unopposed pro-inflammatory response to human cytomegalovirus proteins in vitro are associated with longer survival in very elderly people. *Age (Dordr)* (2013) 35(4):1387–99. doi:10.1007/s11357-012-9425-7
229. Sylwester AW, Mitchell BL, Edgar JB, Taormina C, Pelte C, Ruchti F, et al. Broadly targeted human cytomegalovirus-specific CD4+ and CD8+ T cells dominate the memory compartments of exposed subjects. *J Exp Med* (2005) 202(5):673–85. doi:10.1084/jem.20050882
230. Franceschi C, Bonafe M, Valensin S. Human immunosenescence: the prevailing of innate immunity, the failing of clonotypic immunity, and the filling of immunological space. *Vaccine* (2000) 18(16):1717–20. doi:10.1016/S0264-410X(99)00513-7
231. Akbar AN, Fletcher JM. Memory T cell homeostasis and senescence during aging. *Curr Opin Immunol* (2005) 17(5):480–5. doi:10.1016/j.coi.2005.07.019
232. van Leeuwen EM, Koning JJ, Remmerswaal EB, van Baarle D, van Lier RA, ten Berge IJ. Differential usage of cellular niches by cytomegalovirus versus EBV- and influenza virus-specific CD8+ T cells. *J Immunol* (2006) 177(8):4998–5005. doi:10.4049/jimmunol.177.8.4998
233. Huster KM, Stemmerger C, Gasteiger C, Kastenmuller W, Drexler I, Busch DH. Cutting edge: memory CD8 T cell compartment grows in size with immunological experience but nevertheless can lose function. *J Immunol* (2009) 183(11):6898–902. doi:10.4049/jimmunol.0902454
234. Ferrando-Martinez S, Franco JM, Ruiz-Mateos E, Hernandez A, Ordonez A, Gutierrez E, et al. A reliable and simplified sj/beta-TREC ratio quantification method for human thymic output measurement. *J Immunol Methods* (2010) 352(1–2):111–7. doi:10.1016/j.jim.2009.11.007
235. Radtke F, MacDonald HR, Tacchini-Cottier F. Regulation of innate and adaptive immunity by Notch. *Nat Rev Immunol* (2013) 13(6):427–37. doi:10.1038/nri3445
236. Haugen F, Norheim F, Lian H, Wensaas AJ, Dueland S, Berg O, et al. IL-7 is expressed and secreted by human skeletal muscle cells. *Am J Physiol Cell Physiol* (2010) 298(4):C807–16. doi:10.1152/ajpcell.00094.2009
237. Fry TJ, Mackall CL. The many faces of IL-7: from lymphopoiesis to peripheral T cell maintenance. *J Immunol* (2005) 174(11):6571–6. doi:10.4049/jimmunol.174.11.6571
238. Tchernof A, Despres JP. Pathophysiology of human visceral obesity: an update. *Physiol Rev* (2013) 93(1):359–404. doi:10.1152/physrev.00033.2011
239. Trim W, Turner JE, Thompson D. Parallels in immunometabolic adipose tissue dysfunction with ageing and obesity. *Front Immunol* (2018) 9:169. doi:10.3389/fimmu.2018.00169

240. Nieman DC, Henson DA, Nehlsen-Cannarella SL, Ekkens M, Utter AC, Butterworth DE, et al. Influence of obesity on immune function. *J Am Diet Assoc* (1999) 99(3):294–9. doi:10.1016/S0002-8223(99)00077-2
241. Muezzinler A, Zaineddin AK, Brenner H. Body mass index and leukocyte telomere length in adults: a systematic review and meta-analysis. *Obes Rev* (2014) 15(3):192–201. doi:10.1111/obr.12126
242. van der Weerd K, Dik WA, Schrijver B, Schweitzer DH, Langerak AW, Drexhage HA, et al. Morbidly obese human subjects have increased peripheral blood CD4+ T cells with skewing toward a Treg- and Th2-dominated phenotype. *Diabetes* (2012) 61(2):401–8. doi:10.2337/db11-1065
243. Alam I, Larbi A, Pawelec G. Nutritional status influences peripheral immune cell phenotypes in healthy men in rural Pakistan. *Immun Ageing* (2012) 9(1):16. doi:10.1186/1742-4933-9-16
244. Spielmann G, Johnston CA, O'Connor DP, Foreyt JP, Simpson RJ. Excess body mass is associated with T cell differentiation indicative of immune ageing in children. *Clin Exp Immunol* (2014) 176(2):246–54. doi:10.1111/cei.12267
245. Costanzo AE, Taylor KR, Dutt S, Han PP, Fujioka K, Jameson JM. Obesity impairs gammadelta T cell homeostasis and antiviral function in humans. *PLoS One* (2015) 10(3):e0120918. doi:10.1371/journal.pone.0120918
246. Aroor AR, DeMarco VG. Oxidative stress and obesity: the chicken or the egg? *Diabetes* (2014) 63(7):2216–8. doi:10.2337/db14-0424
247. Gleeson M, Bishop NC, Stensel DJ, Lindley MR, Mastana SS, Nimmo MA. The anti-inflammatory effects of exercise: mechanisms and implications for the prevention and treatment of disease. *Nat Rev Immunol* (2011) 11(9):607–15. doi:10.1038/nri3041
248. Radak Z, Chung HY, Goto S. Systemic adaptation to oxidative challenge induced by regular exercise. *Free Radic Biol Med* (2008) 44(2):153–9. doi:10.1016/j.freeradbiomed.2007.01.029
249. Henson SM, Lanna A, Riddell NE, Franzese O, Macaulay R, Griffiths SJ, et al. p38 signaling inhibits mTORC1-independent autophagy in senescent human CD8(+) T cells. *J Clin Invest* (2014) 124(9):4004–16. doi:10.1172/JCI75051

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Campbell and Turner. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.